

Annex VII

Draft Integrated Report of HABs for the NOWPAP Region

1. Introduction

The purposes of this Integrated Report are to describe HAB problems in the NOWPAP Region and to identify necessary future activities of CEARAC for tackling HAB problems. The information included in this Integrated Report is mainly based on the National Reports submitted by the NOWPAP Members in 2004. Useful supplementary data from other information sources is also used in this Integrated Report.

Figure 1 shows the approximate area of the NOWPAP Region. The Integrated Report covers the part of the NOWPAP Region, which is surrounded by the four countries and their related areas. The reason of the additional areas is that the sea areas outside of the boundary make strong influences to the marine environment of the NOWPAP Region. On the other hand, the Pacific Ocean and the Seto Inland Sea of Japan are not included in this report because WG3 activities are concentrating on problems relevant to the four countries, not to one country of the NOWPAP Members..




Figure 1 Area of the NOWPAP Region

(<http://cearac.nowpap.org/nowpap/coverage.html>)

1.1 Definition of Words

Since each NOWPAP Member has their own definition of HAB, the first Working Group 3 Meeting in Busan, Korea, in October 2003 agreed that WG3 defined HAB. The details are explained as follows. The scientific names of the plankton species are basically described as used in National Reports.

HAB: Proliferation of unicellular plankton, which can cause massive fish or shellfish kills, contaminate seafood with toxins, and alter the aquatic ecosystems in ways that humans perceive as harmful. There are two phenomena, so called Red Tide and Toxin-Producing Plankton.

Red Tide: Water discoloration by vastly increased unicellular plankton that induces deterioration of aquatic ecosystems and occasionally fishery damage.

Toxin-Producing Plankton: Plankton species that produces toxin inside its cell and contaminate fish and shellfish through the food chain.

1.2 Natural Environment of the NOWPAP Region

This section provides a brief overview of the natural environment of the NOWPAP Region, focusing on the three major sea areas, major rivers and ocean currents. Figure 2 shows geographic characteristics of the NOWPAP Region. Compared to Figure 1, Figure 2 includes some outside areas of the boundary of the NOWPAP Region - in B Sea Area and C Sea Area. Data from these areas are included in this Report.




Figure 2 Geographic Characteristics of the NOWPAP Region

1) Sea Area

As shown in Figure 2, A Sea Area, B Sea Area and C Sea Area constitute the major part of the NOWPAP Region’s sea area. Table 1 provides basic information of these three Sea Areas.

Table 1 Basic Information of the Three Seas in the NOWPAP Region

	A Sea Area	B Sea Area	C Sea Area
Surface Area (km ²)	1,300,000	400,000	7,284
Volume (km ³)	1,750,000	17,600	131
Average depth (m)	1,350	44	18
Maximum depth (m)	3,796	100	85

Source: EMECS (2003), Environmental Guidebook on the Enclosed Coastal Seas of the World

A Sea Area is a semi-enclosed sea surrounded by Japan, the Korean Peninsula and Russia. It is connected to the open ocean through several straits. A Sea Area is the largest and deepest sea among the three Sea Areas.

B Sea Area is a semi-enclosed sea bounded by the Chinese mainland on the west, the Korean Peninsula on the east and the East China Sea on the south. The waters of B Sea Area is characterized with yellowish color due to a large amount of yellow silt discharged from large Chinese rivers. The depth of B Sea Area is significantly shallower compared to A Sea Area, having an average depth of only 44 meters.

C Sea Area is the smallest and most enclosed sea within the NOWPAP Region. It is located to the northwest of B Sea Area, and the two sea areas are connected through a relatively wide strait. C Sea Area is even shallower than B Sea Area, with an average depth of 18 meters. C Sea Area functions as an offshore gateway to the capital city Beijing.

2) River

Numerous large and small rivers flow into the three Sea Areas. Table 2 shows some of the major rivers that flow into these three Sea Areas.

Table 2 Major Rivers That Flow into the Three Sea Areas

	River	Members	Catchment Area (km ²)	Flow Rate (m ³ /s)
A Sea Area	Tumangan River	China, Russia	33,800	-
	Nakdong River	Korea	23,817	794
	Tumnin River	Russia	22,400	-
	Ishikari River	Japan	14,330	400
	Shinano River	Japan	11,900	518
B Sea Area	Yangtze River	China	1,807,199	29,000
	Han River	Korea	26,018	1,171
	Kum River	Korea	9,810	841
C Sea Area	Yellow River	China	752,443	1,820
	Haihe River	China	264,617	717
	Liaohe River	China	164,104	302

Source:

NPEC (2003), The State of the Environment of the Northwest Pacific Region.

River Bureau, Ministry of Land, Infrastructure and Transport (2002), River Discharges Year Book of Japan.

Ministry of Construction and Transportation (1998), Discharge Annual Report in Korea.

Some rivers reach to enormous sizes due to mainly their large catchment areas, and they have a significant influence on the NOWPAP Region sea area. Despite their relatively small sizes, B and C Sea Areas receive large amount of river inflow from some of the largest rivers in China such as the Yangtze and Yellow River. Compared among the sea areas above, the rivers in A Sea Area are not so large due to their relatively small catchment areas.


3) Major Oceanographic Currents in the NOWPAP Region

Two strong currents exist in A Sea Area, the Tsushima Warm Current and the Liman Cold Current. The Tsushima Warm Current, a branch of the larger Kuroshio Current, enters A Sea Area from the strait between Japan and Korea and heads toward the northeast. The Liman Cold Current runs along the Eurasian Continent from the north to the south.

The Tsushima Warm Current diverges into three smaller branches upon entering A Sea Area. The first branch goes along the coastline of the Japanese archipelago, the second along the Korean Peninsula then turns and meanders eastward. The third cuts across the center of A Sea Area. Eventually, the major bodies of these currents flow out into the Pacific Ocean or the Sea of Okhotsk through the straits between Hokkaido and Honshu, and Hokkaido and Sakhalin respectively. According to the past records, the Tsushima Warm Current enters A Sea Area and exits into the Pacific Ocean in approximately two months. Some remaining of the current continues to travel northward, slowly cooling down during the travel, and finally exits into the open ocean through the narrow strait between Sakhalin and Russian mainland. Due to shallowness of the strait, part of this current turns around at the mouth of the strait and heads

toward the south along the Eurasian Continent. Finally, it becomes the Liman Current.

The Kuroshio Current also diverges into B Sea Area and C Sea Area as the Yellow Sea Warm Current. Figure 3 is a schematic diagram of the oceanographic currents of the NOWPAP Region.




Prepared based on: Yoon J.H. (1997), Bull. Jpn. Soc. Fish. Oceanogr., 61 (3): 300-303.

Figure 3 Major Oceanographic Currents in the NOWPAP Region

1.3 Social Environment of the NOWPAP Region

1) Demography

The total population in the NOWPAP Region's catchment areas is approximately 560 million, in which approximately 85% are in the Chinese region. Approximately 34 and 47 million people inhabit the Japanese and Korean regions respectively. Only 4.3 million people are in the Russian region. The population density is highest in Korea, followed by China and Japan. The population density in Russia is much smaller than the other NOWPAP Members - two digits less. Figure 4 shows the population and population density in the NOWPAP Region's catchment areas.



Source: NPEC (2003), The State of the Environment of the Northwest Pacific Region

**Figure 4 Populations and Population Densities
 in the NOWPAP Region's Catchment Areas**

2) Aquaculture

Various types of aquaculture are being operated in the NOWPAP Region - cultivating fish, shellfish and seaweeds. Figure 5 shows major aquaculture operating areas in the NOWPAP Region. Aquaculture is widely operated along the coasts of China, Japan and Korea. Although aquaculture in Russia is operated only in limited areas at present, it is expanding. Table 3 shows types of aquaculture conducted in the NOWPAP Region.



Source:

Yoon Y. H. (2001); A summary on the red tide mechanisms of the harmful dinoflagellate, *Cochlodinium polykrikoides* in Korean coastal waters., Bull. Plankton Soc. Japan, 48 (2): 113-120.

Matsuoka K. (2004); Present status in study on a harmful unarmored dinoflagellate *Cochlodinium polykrikoides* Margalef., Bull. Plankton Soc. Japan, 51 (1): 38-45.

Geological Institute, China Scientific Academy (1999); Chinese national atlas of natural resources

Figure 5 Major Aquaculture Areas in the NOWPAP Region

Table 3 Types of Aquaculture Conducted in the NOWPAP Region

	Location	Type of Aquaculture
China	Coast of Bohai Sea, Shandong Peninsula, Liaodong Peninsula	Tiger prawns, Seaweeds, etc.
Japan	North coast of Kyushu West coast of Hokkaido	Amberjack, Red seabream Scallop
Korea	West and south coast	Bastard Halibut, Amberjack, Rockfish
Russia	South coast of Sakhalin, Peter the Great Bay	Scallop, Seaweeds, Mussel, Cucumaria

2. Situation of HAB Occurrence

2.1 Present Situation of HAB in the NOWPAP Region

In this chapter, the status of HAB in the NOWPAP Region is summarized. Information on red tides and toxin-producing plankton is presented separately.

1) Red Tides

Table 4 summarizes the status of red tide events in the NOWPAP Region. Approximate locations of red tide events are shown in Figure 6(See p.14). Despite the fact that the HAB monitoring activity does not cover all the coastal areas in the NOWPAP Region (see Section 3.1), red tide events have been constantly recorded along the coastal areas with yearly and spatial variations. Intensive fishery and aquaculture areas tend to have many records of red tide occurrences.

So far 75 species have been recorded in the NOWPAP Region (Table 5). Three flagellate species (*Heterosigma akashiwo*, *Noctiluca scintillans*, *Prorocentrum minimum*) and one diatom species (*Skeletonema costatum*) have been recorded in the coastal waters of all the NOWPAP Members. All the three flagellate species above have caused extensive damage to local fisheries. Other common and damage-causing species include *Gymnodinium mikimotoi*, *G. sanguineum* and *Prorocentrum micans* (all flagellates). In recent years *Cochlodinium polykrikoides* has caused serious damage to fisheries in Japan and Korea.

The extent of red tides within the NOWPAP Region is usually limited to less than 100km² in Japanese, Korean and Russian waters. Blooms in Chinese waters, however, often extend over a 100 km². More than 50% of blooms in data between 1990 and 2004 were larger than 100 km², and approximately 25% of them were larger than 1,000 km² (refer to table 4). One of the reasons of this difference in data results occurred between China and the other NOWPAP Members may be because of their different data sources.. In China, bloom size was mostly identified through aerial survey, whereas the other NOWPAP Members adopted data mainly through vessels.

Red tides were most frequent from spring to summer in the NOWPAP Region. Figure 7 shows monthly patterns of red tide events in the NOWPAP Region. The peak season in China was from June to August. The peak in Japan was in April, June and July. In Korea, there was a prominent peak in August. In Russia, the peak appeared in June and July. The dominant red tide species during the peak months were as follows:

China; *Noctiluca scintillans* (June and July)

Japan; *Noctiluca scintillans* (April), *Heterosigma akashiwo* (June), *Gymnodinium mikimotoi* (July)

Korea; *Cochlodinium polykrikoides* (August)

Russia; *Noctiluca scintillans* and *Heterosigma akashiwo* (June)

All the species above are known to cause damage to fisheries.

Most red tide events in the NOWPAP Region lasted for about one week. In some rare cases, however, red tides lasted for one to two months (e.g. *C. polykrikoides* bloom lasted for 62 days in Korea in 2003).

Several mitigation measures have been developed or are under development to counteract red tide blooms. Clay spraying is one of the common methods employed in the NOWPAP Region.

Table 4 Summary of Red Tide Events in the NOWPAP Region

	China (Bohai and Yellow Sea)	Japan (Data from Kyushu region unless stated (1998-2002))	Korea (1999-2003 unless stated)	Russia (1992-2003 unless stated)*1
Number of events	84 red tide events from 1990-2004.	150 red tide events recorded. 19 events were harmful.	304 red tide events recorded.	23 red tide events recorded. All events were harmless and caused no damage.
Causative species	Refer to Table 5	Refer to Table 5 (includes Honshu region)	Refer to Table 5	Refer to Table 5
Cell density	Maximum cell density recorded for the following major red tide species: <i>Noctiluca scintillans</i> (49,000 cells/ml) <i>Skeletonema costatum</i> (72,000 cells/ml) <i>Ceratium furca</i> (1,250 cells/ml) <i>Gymnodinium</i> sp. (300,000 cells/ml)	<i>Gymnodinium mikimotoi</i> recorded the highest density at 117,980 cells/ml.	Each year <i>Cochlodinium polykrioides</i> recorded the highest cell density. Maximum cell density was recorded in 2003 at 48,000 cells/ml.	<i>Eutreptiella gymnastica</i> recorded the highest density at 30,900 cells/ml.
Location of occurrence	Mainly along the coast of C Sea Area (Refer to Figure 6)	Mainly along the coast of northern Kyushu (Refer to Figure 6: includes Honshu region)	Along the entire coast except the northeast coast (Refer to Figure 6)	Some areas in Peter the Great Bay (Refer to Figure 6)
Size of bloom	Data from 1990-2004 <10km ² : 18% 10-100km ² : 29% 100-1,000km ² : 30% >1,000km ² : 23% Affected area generally larger in C Sea Area compared to B Sea Area.*2	<1km ² : 51% 1-100km ² : 48% >100km ² : 1%	<1km ² : 56% 1-100km ² : 19% >100km ² : 24% Large blooms were mostly by <i>C. polykrioides</i> .	<i>Noctiluca scintillans</i> and <i>Prorocentrum minimum</i> blooms exceeded 1km ² .
Duration	Most red tides lasted less than a week. However, <i>Ceratium furca</i> bloom lasted for 40 days in 1998. <i>Eucampia zodiacus</i> and <i>Chaetoceros socialie</i> bloom lasted for 20 days.	Although there were some variations, red tide events tended to last around 1 week. 18 out of 150 events lasted over 20 days.	Most red tide lasted for less than 10 days except for <i>C. polykrioides</i> , which continued for 1 – 2 months.	<i>N. scintillans</i> and <i>Oxyrrhis marina</i> blooms lasted more than 20 days.
Seasonal pattern	Most frequent in July and August (1990-2004). Refer to Figure 7 for more details.	High frequency of red tide between April – September. Most frequent in June and July. Refer to Figure 7 for more details.	Red tides recorded from January – November. Most frequent in August. Refer to Figure 7 for more details.	Mostly observed between March – September. Most frequent in June and July. Refer to Figure 7 for more details.
Damage	Mass mortality of fish and shellfish by: <i>Ceratium furca</i> , <i>Exuviaella cordata</i> , <i>Gymnodinium</i> sp., <i>G. sanguineum</i> , <i>N. scintillans</i> , <i>Prorocentrum</i> sp. Most serious damage recorded in 1989 by <i>Gymnodinium</i> sp. in Bohai Bay (economic loss of US\$ 38 million).	Mass mortality of fish and shellfish by: <i>Heterosigma akashiwo</i> , <i>Heterocapsa circularisquama</i> , <i>G. mikimotoi</i> , <i>C. polykrioides</i> , <i>N. scintillans</i> Most serious damage recorded in 1999 by <i>C. polykrioides</i> (economic loss of US\$ 7 million)	<i>C. polykrioides</i> has caused damage to fisheries for most years since 1993. Economic loss of US\$ 95 million in 1995 and US\$ 19 million in 2003.	No damage recorded.
Mitigation measures	Regular monitoring (more details in Ch.3) Preventive measures: Effluent control (implementation of Blue Sea Action Plan), improvement of sewage system, public education Reactive measure: Aeration of seawater and fish-pen sinking in fish farms, clay spraying	Regular monitoring (more details in Ch.3) Preventive measures: Effluent control, improvement of sewage system, public education Reactive measures: Clay spraying	Regular monitoring (more details in Ch.3) Deployment of Automatic HAB Alarm System in aquaculture farms. Reactive measures: Clay spraying, Electrolytic Clay Dispenser (ECD)	No mitigation measures employed.

*1: There are no regular red tide monitoring programs in Russia. The presented data is referred from ad hoc monitoring or research conducted by the IMB FEB RAS in 1992 – 2002.

*2: Observation was mainly conducted through aerial survey.

Table 5(1) Red Tide Species Recorded in the NOWPAP Region

Class	Genus and Species	China	Japan	Korea	Russia
Bacillariophyceae	<i>Asterionella</i> sp.		✓		
	<i>Chaetoceros curvisetum</i>		✓		
	<i>Chaetoceros socialie</i>	✓			
	<i>Chaetoceros</i> sp.		✓	✓	
	<i>Coscinodiscus asteromphalus</i>	✓			
	<i>Coscinodiscus gigas</i>			✓	
	<i>Coscinodiscus</i> sp.			✓	
	<i>Ditylum brightwellii</i>				✓
	<i>Eucampia zodiacus</i>	✓		✓	
	<i>Eucampia</i> sp.			✓	
	<i>Leptocylindrus danicus</i>	✓	✓	✓	
	<i>Leptocylindrus</i> sp.		✓		
	<i>Navicula</i> sp.	✓			
	<i>Neodelphineis pelagica</i>		✓		
	<i>Nitzschia</i> sp.		✓	✓	
	<i>Pseudo-nitzschia calliantha</i>				✓
	<i>Pseudo-nitzschia multiseriis</i>				✓
	<i>Pseudo-nitzschia pseudodelicatissima</i>				✓
	<i>Pseudo-nitzschia pungens</i> ^{*1}			✓	✓
	<i>Pseudo-nitzschia</i> sp.			✓	
	<i>Rhizosolenia delicatula</i>			✓	
	<i>Rhizosolenia fragilissima</i>			✓	
	<i>Rhizosolenia setigera</i>			✓	
	<i>Rhizosolenia</i> sp.	✓	✓	✓	
	<i>Skeletonema costatum</i>	✓	✓	✓	✓
	<i>Skeletonema</i> sp.			✓	
	<i>Thalassiosira decipiens</i>			✓	
	<i>Thalassiosira rotula</i>			✓	
<i>Thalassiosira</i> sp.			✓		
Cyanophyceae	<i>Microcystis viridis</i>			✓	
Dinophyceae	<i>Alexandrium catenella</i>	✓	✓		
	<i>Alexandrium fraterculus</i>		✓		
	<i>Alexandrium</i> sp.			✓	
	<i>Ceratium furca</i>	✓	✓		
	<i>Ceratium fusus</i>			✓	
	<i>Ceratium</i> sp.			✓	
	<i>Cochlodinium polykrikoides</i>		✓	✓	
	<i>Cochlodinium</i> sp.		✓		
	<i>Exuviaella cordata</i>	✓			
	<i>Exuviaella marina</i>	✓			
	<i>Dinophysis ovata</i>	✓			

Table 5(2) Red Tide Species Recorded in the NOWPAP Region

Class	Genus and Species	China	Japan	Korea	Russia
Dinophyceae	<i>Gonyaulax spinifera</i>	✓			
	<i>Gymnodinium mikimotoi</i>	✓	✓	✓	
	<i>Gymnodinium sanguineum</i>	✓	✓	✓	
	<i>Gymnodinium sp.</i>			✓	
	<i>Gyrodinium sp.</i>	✓	✓		
	<i>Heterocapsa circularisquama</i>		✓		
	<i>Heterocapsa sp.</i>			✓	
	<i>Heterocapsa triquetra</i>			✓	
	<i>Noctiluca scintillans</i> ^{*2}	✓	✓	✓	✓
	<i>Oxyrrhis marina</i>				✓
	<i>Prorocentrum balticum</i>		✓		
	<i>Prorocentrum dentatum</i>		✓	✓	
	<i>Prorocentrum micans</i>	✓	✓	✓	
	<i>Prorocentrum minimum</i>	✓	✓	✓	✓
	<i>Prorocentrum sigmoides</i>		✓		
	<i>Prorocentrum triestinum</i>		✓	✓	
<i>Prorocentrum sp.</i>			✓		
Raphidophyceae	<i>Chattonella antiqua</i>	✓	✓		
	<i>Chattonella globosa</i>				✓
	<i>Chattonella marina</i>	✓	✓		
	<i>Fibrocapsa japonica</i>		✓		
	<i>Heterosigma akashiwo</i> ^{*3}	✓	✓	✓	✓
Chrysophyceae	<i>Dictyocha fibula</i>			✓	
Eugrenophyceae	<i>Eutreptia lanowii</i>				✓
	<i>Eutreptiella gymnastica</i>		✓	✓	✓
	<i>Eutreptiella sp.</i>			✓	
Haptophyceae	<i>Phaeocystis sp.</i>	✓			
Cryptophyceae	<i>Chroomonas marina</i>			✓	
	<i>Chroomonas salina</i>			✓	
	<i>Cryptomonas acuta</i>			✓	
	<i>Cryptomonas sp.</i>			✓	
Prasinophyceae	<i>Pyramimonas sp.</i>		✓		
Ciliate	<i>Mesodinium rubrum</i>	✓	✓	✓	
	<i>Tontonia sp.</i>		✓		

*1: *Nitzschia pungens* is the synonym of *Pseudo-nitzschia pungens*. In this Report, *N. pungens* is referred to as *P. pungens*


*2: *Noctiluca scintillans* is the sole species of the genus. Therefore, *Noctiluca sp.* is included into *N. scintillans*.

*3: *Heterosigma akashiwo* is the sole species of the genus. Therefore, *Heterosigma sp.* is included into *H. akashiwo*.



Figure 6 Locations of Red Tides in the NOWPAP Region in 1999-2002

The number of red tides	
China	38
Japan	148
Korea	259
Russia	12



Note: Surveyed periods and sample numbers differ among the NOWPAP Members

Figure 7 Seasonal Patterns of Red Tide Occurrences in the NOWPAP Region

2) Toxin-Producing Plankton and Shellfish Poison

Table 6 shows the status of toxin-producing plankton and shellfish poison in the NOWPAP Region. In this Report, toxin-producing species are separated into PSP-, DSP- or ASP-inducing species instead of its taxonomic classification.

(1) Main toxin-producing species

A total of 20 toxin-producing plankton species have been recorded in the NOWPAP Region (Table 7). Six species were PSP-inducing species. All PSP species except *Gymnodinium catenatum* belonged to the genus *Alexandrium*. The most common recorded PSP species in the NOWPAP Region was *A. tamarense*.

Nine species out of the ten DSP species recorded in the NOWPAP Region belonged to the genus *Dinophysis*. The rest was *Exuviaella marina*, which was recorded only in China. Among the *Dinophysis* species, *D. fortii* and *D. acuminata* were recorded in all of the NOWPAP Members.

Damage from ASP has not been recorded yet in the NOWPAP Region, although ASP-inducing *Pseudo-nitzschia* species were recorded in Russia and Korea.

(2) Affected areas

PSP has been recorded in the Shangdong Peninsula and Lianyungang Area in China (Figure 8). Areas affected by PSP in Japan are concentrated in the western Japan (Kyushu and Chugoku region) and Tohoku region (Aomori Prefecture) as shown in Figure 9. In Korea, PSP has recently affected shellfish harvesting areas on the southeastern coast. Russia has not been affected by PSP as yet.

DSP species has been recorded in the Shangdong Peninsula, Lianyungang Area and C Sea Area in China. A *Dinophysis ovata* blooms were recorded over an area of 5,000 km² in C Sea Area in 1998. Areas affected by DSP in Japan are mainly in Hokkaido region, Tohoku region and Chugoku region. In Korea, three *Dinophysis* species were recorded on the southeastern coast in 2002 and 2003, however, it is uncertain whether there was any damage or not by the species. Russia has not been affected by DSP as yet.

In Russia, observation of PSP-, DSP- or ASP-inducing species are conducted mainly in aquaculture areas. Figure 10 – 12 shows the results of these observations. Although incidents of shellfish poison have not been reported in these aquaculture areas as yet, the presence of toxin-producing plankton has been recorded constantly.

(3) Damage

In China, more than 600 people have suffered from shellfish poison since 1967, in which thirty fatalities have resulted from PSP. In Japan, approximately 900 people have suffered from PSP or DSP since 1976, including several deaths from PSP. In Korea, shellfish harvesting was banned on the southeastern coast in 2002 (April – May) and 2003 (April – June) due to *A. tamarense*.

(4) Mitigation measures

China, Japan and Korea are conducting policies to prevent and lessen damage by toxins of shellfish to people. These countries monitor toxin level of the shellfish at harvest areas. When the toxin level exceeds the quarantine limit set by each country, they advise that fishery markets stop shipping or ban harvest of shellfish in some period.

Table 6 Status of Toxin-Producing Plankton and Shellfish Poison in the NOWPAP

	Region			
	China	Japan	Korea	Russia
Main toxin-producing species	<i>Alexandrium catenella</i> , <i>Dinophysis fortii</i> , <i>D. acuminata</i> , <i>D. ovata</i> and <i>Exuviaella marina</i> (Refer to Table 7)	<i>Alexandrium tamarense</i> , <i>A. catenella</i> , <i>A. tamiyavanichii</i> , <i>Gymnodinium catenatum</i> , <i>Dinophysis fortii</i> , <i>D. acuminata</i> , <i>D. caudate</i> , <i>D. intundibrus</i> , <i>D. mitra</i> and <i>D. rotundata</i> (Refer to Table 7)	<i>Alexandrium tamarense</i> , <i>Dinophysis fortii</i> , <i>D. acuminata</i> , <i>D. caudate</i> , <i>D. rotundata</i> and <i>Pseudo-nitzschia pungens</i> (Refer to Table 7)	<i>Alexandrium tamarense</i> , <i>A. acatenella</i> , <i>A. pseudogonyaulax</i> , <i>Dinophysis fortii</i> , <i>D. acuminata</i> , <i>D. acuta</i> , <i>D. norvegica</i> , <i>D. rotundata</i> , <i>Pseudo-nitzschia calliantha</i> , <i>P. multiseriata</i> , <i>P. pseudodelicatissima</i> and <i>P. pungens</i> (Refer to Table 7)
Affected species	Information available only for the southern region of China (outside the NOWPAP Region). PSP: Marine snail (<i>Nussarius succinstus</i>), Clam (<i>Soletellina diphos</i> , <i>Ruditapes phillipenensis</i> , <i>Pinna pectinata</i>), Mussel (<i>Perna viridis</i>)	PSP: Mediterranean blue mussel, Japanese oyster, noble scallop DSP: Mediterranean blue mussel, Japanese scallop	Information N/A	No shellfish poison reported.
Affected area	Shangdong Peninsula, Lianyungang area and C Sea Area (see Figure 8)	Mainly in Hokkaido region, Tohoku region and Chugoku region (see Figure 9)	Southeast coast (Gosung, Tongyoung, Jinhaeman)	No shellfish poison reported. Cell density of potential causative species recorded in certain areas (see Fig.10-12)
Damage	More than 600 people have suffered from shellfish poison since 1967. 30 fatalities from PSP across the nation.	Approximately 900 people have suffered from PSP or DSP since 1976 including several deaths from PSP. No fatalities since 1980.	Stoppage of shellfish harvest in 2002 and 2003 in the southeast coast due to PSP.	No damage was recorded.
Mitigation measures	Some SOA laboratories and local fishery environmental laboratories conduct monitoring of toxin-producing plankton and shellfish poison.	Regular monitoring of main toxin-producing species and toxin test of harvested shellfish. Shipping is voluntarily stopped if the toxicity exceeds the Fishery Agency standard. (Voluntary Control) PSP: 20 cases of voluntary control in 1978-1999. Most cases lasted for 2-4 months. DSP: 64 cases of voluntary control in 1978-1999. Duration of DSP was generally longer than PSP. Some cases lasted for over 5 months.	Regular monitoring of <i>Alexandrium</i> sp. and toxin test of harvested shellfish. Harvest is stopped when the toxin level exceeds the quarantine limit.	No mitigation measures or monitoring.


Table 7 Toxin-Producing Plankton Species Recorded in the NOWPAP Region

	Species name	China	Japan	Korea	Russia
PSP	<i>Alexandrium acatenella</i>				✓
	<i>Alexandrium tamarense</i>		✓	✓	✓
	<i>Alexandrium catenella</i>	✓	✓		
	<i>Alexandrium pseudogonyaulax</i>				✓
	<i>Alexandrium tamiyavanichii</i>		✓		
	<i>Gymnodinium catenatum</i>		✓		
DSP	<i>Dinophysis fortii</i>	✓	✓	✓	✓
	<i>Dinophysis acuminata</i>	✓	✓	✓	✓
	<i>Dinophysis acuta</i>				✓
	<i>Dinophysis caudata</i>		✓		
	<i>Dinophysis infundibrus</i>		✓		
	<i>Dinophysis mitra</i>		✓		
	<i>Dinophysis norvegica</i>				✓
	<i>Dinophysis ovata</i>	✓			
	<i>Dinophysis rotundata</i>		✓	✓	✓
	<i>Exuviaella marina</i>	✓			
ASP ^{*1}	<i>Pseudo-nitzschia calliantha</i>				✓
	<i>Pseudo-nitzschia multiseriis</i>				✓
	<i>Pseudo-nitzschia pseudodelicatissima</i>				✓
	<i>Pseudo-nitzschia pungens</i>			✓	✓

*1:Damage from ASP has not been recorded yet in the NOWPAP Region, although ASP inducing *Pseudo-nitzschia* species were recorded in Russia and Korea according to the National Report. ASP-inducing species probably exist in China and Japan as well, but it has not being recorded due to different monitoring methods. ASP in the NOWPAP Region should be investigated in the future.



Figure 8 Areas where Shellfish Toxin have been Recorded in Coastal China



Source: Japan Fisheries Resource Conservation Association (JFRCA), 'Monitoring Report on Shellfish Poison in Japanese Fishery Products', 1999-2000

Figure 9 Areas that Experienced Voluntary Control due to PSP and DSP Contamination in Japan (1978-1999)




Figure 10 Dates of Occurrences and Maximum Cell Densities of *Alexandrium* Species in Russian Coastal Waters in 1992-2002




Figure 11 Dates of Occurrences and Maximum Cell Densities of *Dinophysis* Species in Russian Coastal Waters in 1992-2002




Figure 12 Dates of Occurrences and Maximum Cell Densities of *Pseudo-nitzschia* Species in Russian Coastal Waters in 1992-2002

2.2 Common issues on HAB in the NOWPAP Region

1) Severe fishery damage caused by *Cochlodinium polykrikoides*

Red tides have often resulted in large mortality of fishery resources and huge economic loss to the fisheries in the NOWPAP Region. They often occur in semi-enclosed areas such as inlets and embayments where aquaculture is often operated. Although various species are known to cause red tides, in recent years *C. polykrikoides* has caused the most serious damage to the fisheries in Japan and Korea. For example in 1999, approximately US\$ 7 million worth of fishery damage was recorded in Imari Bay, Kyushu, Japan. Even greater economic losses were recorded in Korea in 1995 and 2003, worth at approximately US\$ 95 million and US\$ 19 million respectively. Based on National Reports and recent research papers, the locations of *C. polykrikoides* blooms in the Japanese and Korean regions are plotted in Figure 13.

To prevent or lessen future damage from *C. polykrikoides*, various researches have been conducted to understand the ecology of this species. Several researches have focused on transportation scheme of this species. Miyahara et al. (2005) traced the movement of *C. polykrikoides* blooms that occurred along the A Sea Area coast of Chugoku region in 2003, by detecting the chlorophyll-a concentration taken in the satellite images of the area (field measurements verified that the high chlorophyll-a concentration in the satellite images was formulated predominantly by *C. polykrikoides*). Figure 14 shows how the *C. polykrikoides* blooms moved along the coast of Chugoku region. Miyahara et al. concluded that this particular *C. polykrikoides* bloom was most likely transported to the coast of Chugoku region through the Tsushima Warm Current.

Kim et al. (2004) have studied the impact of water temperature, salinity and irradiance on the growth rate of *C. polykrikoides*. The highest growth rate was recorded when the water temperature was 25°C, salinity 34 and irradiance >90 μ mol/m²/s. these may explain the appropriate conditions of the *C. polykrikoides* blooms recorded in the Japanese (Kyushu region) and Korean regions. All the *C. polykrikoides* blooms occurred between August and October in these areas when the water temperature was close to the value above. However, the optimum growth conditions of *C. polykrikoides* should be studied further through collection of field data.




Source:

Yoon Y. H. (2001); A summary on the red tide mechanisms of the harmful dinoflagellate, *Cochlodinium polykrikoides* in Korean coastal waters., Bull. Plankton Soc. Japan, 48 (2): 113-120.

Matsuoka K. (2004); Present status in study on a harmful unarmored dinoflagellate *Cochlodinium polykrikoides* Margalef., Bull. Plankton Soc. Japan, 51 (1): 38-45.

Figure 13 Locations of *C. polykrikoides* Blooms in Japan and Korea



Note: The movement of *C. polykrikoides* blooms along the coast of Chugoku region is clearly seen with green color from September 4th to 7th. The spread of primary production on September 16th and 18th is thought to be caused by the typhoon on September 12th.

Source: Miyahara et al. (2005): A harmful bloom of *Cochlodinium polykrikoides* Margalef (Dinophyceae) in the coastal area of San-in, western part of the Japan Sea, in September 2003, Bull. Plankton Soc. Japan, 52(1), 11-18.

Figure 14 Movement of *C. polykrikoides* Blooms along the Coast of Chugoku Region in A Sea Area

2) Threats of DSP and PSP

Shellfish poison is a common threat in the NOWPAP Region. In China, more than 600 people have suffered from shellfish poison since 1967, in which 30 cases were fatal. The majority of these fatalities were from PSP. In Japan, approximately 900 people have suffered from shellfish poison of DSP and PSP since 1976. In Korea, shipping of shellfish was temporarily suspended in 2002 and 2003 due to PSP. Although there have been no reports of shellfish poison incidents in Russia as yet, the presence of various toxin-producing species have been recorded in Russian waters. Shellfish poison in Russia could become a major threat in the future, especially in parallel with the expansion of the aquaculture industry.

3. Information of HAB Monitoring

3.1 Monitoring Activities in the NOWPAP Region

Table 8 summarizes the current status of HAB monitoring in the NOWPAP Region. The locations of monitored areas are shown in Figure 15 and 16.

1) Monitoring of red tide

Apart from Russia, all the NOWPAP Members have a regular red tide monitoring program, though monitoring efforts and methods vary with the NOWPAP Members. In China and Japan, red tide monitoring sites are distributed sporadically over the country, usually conducted in areas with high aquaculture activities. In Korea, red tide monitoring sites are distributed densely over the entire coast. Regular monitoring in Russia has not been established yet partly due to the small number of aquaculture farms in the Far East coast. However, realizing the recent increases of red tide events and its potential negative effects to fisheries, IMB FEB RAS has conducted several red tide (monitoring and) researches on an ad hoc basis.

Red tide monitoring in China, Japan and Korea are mainly conducted by fisheries research organizations. Other national institutes also provide valuable information on red tides through aerial surveys, satellite data, and so on. In case of a significant red tide event, various institutes collaborate to conduct trace monitoring and implement effective countermeasures. Korea especially has a well established inter-organization cooperation scheme for such cases through the NFRDI HAB Emergency Center.

2) Monitoring of toxin-producing Plankton

Monitoring of toxin-producing plankton is conducted in China, Japan and Korea, usually by fisheries research organizations. In Japan, the monitoring is conducted in selected shellfish-producing areas.

In Japan and Korea, monitoring is usually focused on certain target species. However, each fisheries research organizations set their targets on different species. In Japan, *Alexandrium* species and *Gymnodinium catenatum* are usually monitored for PSP, and *Dinophysis* species for DSP. In Korea, *Alexandrium tamarense* is monitored in the southeastern region near aquaculture farms.

3) Monitoring of shellfish poison

Monitoring of shellfish poison is conducted in China, Japan and Korea, usually by fisheries research organizations. In Japan and Korea, this type of monitoring is conducted in

shellfish-producing areas.

All the NOWPAP Members have quarantine limits for harvested shellfish. When the toxin level exceeds the limit, shipping or harvesting of shellfish is stopped until the toxin values return to acceptable levels. The limit for PSP established in China, Korea and Russia is $80 \mu\text{g}$ (STX eq.)/100g of whole meat. Japan applies Mouse Unit (MU) for expressing the toxic level. Japanese standards are—PSP: 4MU/g, DSP: 0.05MU/g. Some researchers mention a report in which it is said that 1MU/g is equivalent to approximately $20 \mu\text{g}$ (STX eq.)/100g.

3.2 Common Issues on Monitoring Activities in the NOWPAP Region

Although HAB monitoring is conducted by all the NOWPAP Members, there are some variations in their monitoring methods and the amount of monitoring efforts among the NOWPAP Members. Such variations have resulted from differences of HAB problems, and restrictions of personnel, technology and finance. For example, Russia does not have a strong demand for HAB monitoring as Japan and Korea, since Russian aquaculture activities is still relatively small.

Another thing which makes the situation more complicated is local variations of monitoring schemes, which is apparent in China and Japan. For example in Japan, the method of HAB monitoring varies with each prefectural fisheries laboratory. This variation happens because they conduct HAB monitoring in accordance to indigenous species and budget for monitoring. As a result, a consistent methodology for HAB monitoring has not been established nationwide. Furthermore, monitoring could be stopped if prefectural fisheries laboratories can not obtain budget for HAB monitoring.

Table 8 Status of HAB Monitoring in the NOWPAP Region

		China		Japan			Korea		Russia
Red tide (regular monitoring)	Major implementing organization	Branch office of SOA	SEPA Department of Agriculture Fishery environmental laboratories of local government	Fishery laboratories of prefectural governments	Kyushu Fisheries Coordination Office	Japan Coast Guard	NFRDI NFRDI fisheries extension service center	National Maritime Police Agency (NMPA)	No regular government monitoring program. However, IMB FEB RAS conduct observations on ad hoc basis.
	Method	Vessel monitoring Satellite remote sensing Aerial monitoring	Information N/A	Temperature, salinity, chlorophyll-a, nutrients, cell density monitored at fixed points (some labs do not monitor all these parameters)	Water color (visual observation) and water temperature (infrared sensor) monitored through aerial survey.	Information N/A	Cell density of <i>C. polykrikoides</i> . Precaution and warnings issued when <i>C. polykrikoides</i> cell density exceed 300 cells/ml and 1,000 cells/ml respectively.	Aerial survey	Information N/A
	Location	4 monitoring sites in Yellow and Bohai Sea. See Figure 15 for location.	Information N/A	Usually limited to small area such as in enclosed bays. See Figure 15 for the monitored site.	4 flight routes over the Kyushu coastal area.	Offshore areas	169 stations. See Figure 15 for location.	Information N/A	Coastal waters of Primorye and South Sakhalin Island.
	Frequency	Information N/A	Information N/A	Differ with the laboratories. Mainly during spring to summer.	6-8 flights during June-October.	Information N/A	February – November	Information N/A	Ad hoc basis
Red tide (trace monitoring)		After the initiation of red tide, fishery environmental laboratories conduct plankton sampling and when necessary continue tracking. SOA also participate in tracking when required.		After the initiation of red tide, fishery laboratories conduct plankton sampling and when necessary continue tracking.			After the initiation of red tide, HAB Emergency Center in NFRDI collect relevant information to predict the future movement of red tide. The information is then disseminated to fishermen and relevant organizations.		Trace monitoring not conducted.
Toxin-producing plankton	Implementing organization	Some SOA laboratories and local fishery environmental laboratories. Monitoring network under construction.		Fishery laboratories of prefectural governments			NFRDI and Regional Maritime Affairs and Fisheries Office		No official regular monitoring program. However, IMB FEB RAS and SakhNIRO conduct observations at an ad hoc basis.
	Method	Information N/A		Cell density of <i>Alexandrium</i> species and <i>Gymnodinium catenatum</i> are usually monitored for PSP, and <i>Dinophysis</i> species for DSP. However, the target species may differ with the laboratories.			Cell density of <i>A. tamarensis</i> is regularly monitored.		Cell density of certain toxin-producing plankton studied.
	Location	Information N/A		Usually limited to small area such as in enclosed bays. See Figure 16 for the monitored area.			Near the shellfish farms in the southeast coast.		Coastal waters of Primorye and South Sakhalin Island.
	Frequency	Information N/A		Differ with the laboratories.			Information N/A		Ad hoc basis
Shellfish poison	Implementing organization	Some SOA laboratories and local fishery environmental laboratories. Monitoring network under construction.		Fishery laboratories of prefectural governments			NFRDI and Regional Maritime Affairs and Fisheries Office		Monitoring not conducted.
	Method	Information N/A		Measurement of toxin level in the midgut gland.			Measurement of toxin level in the meat or midgut gland.		-
	Location	Information N/A		Fish landing ports. See Figure 16 for the monitored area.			Shellfish farms in the western and southern coastal area. Over 100 stations. See Figure 16 for the monitored area.		-
	Frequency	Varies with the local harvest season.		At least monthly during the harvest season. Frequency increases to weekly if high risk of poison is suspected.			At least more than once a month. Frequency increases when toxin is detected in the shellfish.		-
	Shipping harvest stoppage	Stoppage of harvesting and shipping when PSP toxin level exceeds the Department of Agriculture standard (80 μg/100g of whole meat). DSP toxin level must be non-detectable.		Voluntary stoppage of shipping when toxin level exceeds the Fishery Agency standard (PSP: 4MU/g, DSP: 0.05MU/g). Shipping can recommence when toxicity level remain below the standard for 2 weeks.			Stoppage of harvesting when PSP toxin level exceeds 80 μg/100g meat.		Maximum permissible level. PSP: 80 μg/100g wet mollusk tissue. DSP: No detection of ocaidaic acid.



Note 1: Blue plots show the locations of monitoring organizations in Japan. Actual monitoring sites are usually in the adjacent coastal waters of the monitoring organizations.

Note 2: Only SOA monitoring sites are shown for China.

**Figure 15 Monitoring Sites of Red Tides in the NOWPAP Region as of 2002
(including trace monitoring)**



Note 1: Blue plots show the locations of monitoring organizations. Actual monitoring sites are usually in the adjacent coastal waters of the monitoring organizations.

Note 2: Monitoring of shellfish poison is conducted only in some of the areas bracketed in red, usually by local fishery laboratories.

**Figure 16 Regular Monitoring Areas of Toxin-producing Plankton and Shellfish Poison
in the NOWPAP Region as of 2002**

4. Researches and Studies to Cope with HAB

Table 9 shows main HAB researches and studies conducted in the NOWPAP Region, categorized into the mechanism of HAB occurrences, toxicity analysis, taxonomy and mitigation measures.

The bloom mechanisms of harmful species were investigated in relation to various physical, chemical and biological environmental parameters. Target species include *Alexandrium* spp., *Gymnodinium* sp. and *C. polykrikoides*. Some researches have focused on interspecific relationships among plankton, bacteria and virus species as a key to initiate or extinguish the population of harmful plankton species.

Toxicity analysis is one of the hot topics. The effectiveness of various new assay or bioassay techniques are being tested to improve their detection abilities. The toxicity of various harmful species, including intraspecies variation is also studied.

Recent studies on plankton taxonomy are characterized with a molecular biological approach. It has been employed for species identification, intraspecies genetic variation, and so on.

Possible new mitigation measures are constantly being researched in the NOWPAP Region. Physical control of HAB through clay spraying is a well studied method and has been already implemented in some areas. However, its environmental impact is still under concern. The use of surfactant has also been considered by some studies. Biological control of HAB has been recently considered as an effective option by some researchers. Biological methods may control HABs by introducing organisms that graze (e.g. zooplankton, other micro algae, etc) or infect (virus, bacteria) the target plankton species, though the ecological impact should be examined carefully.

Forecast of HAB is another major research topic in the NOWPAP Region. The use of satellite remote sensing is considered as the most effective tool for forecasting the appearance of HABs. Neural network techniques and numerical simulation models are also being studied for predicting the occurrence and movement of blooms.

Table 9 Major HAB Researches and Studies Conducted in the NOWPAP Region

Category	China	Japan	Korea	Russia
Mechanism of HAB occurrence	<ul style="list-style-type: none"> ➤ Relationship of nutrient level with HAB ➤ Relationship of zooplankton community structure with HAB ➤ Bloom mechanism of <i>A. tamarense</i> ➤ Relationship of macronutrients with HAB ➤ Relationship of <i>Alexandrium</i> sp. growth with bacteria ➤ Relationship of <i>A. tamarense</i> growth with Fe and Mn 	<ul style="list-style-type: none"> ➤ Bloom mechanism of PSP inducing species (<i>Alexandrium</i> spp. and <i>Gymnodinium. catenatum</i>) ➤ Relationship of bacteria/viruses with red tide senescence ➤ Relationship of water temp., salinity and irradiance with <i>Cochlodinium. polykrikoides</i> growth 	<ul style="list-style-type: none"> ➤ Relationship of zooplankton community structure with <i>C. polykrikoides</i> bloom ➤ Relationship of physico-chemical factors (water temp., salinity, irradiance and nutrient) with <i>C. polykrikoides</i> bloom 	<ul style="list-style-type: none"> ➤ Bloom mechanism of diatom <i>Chaetoceros salsaugineus</i> and <i>Oxyrrhis marina</i> ➤ Relationship of nutrient level, stratification and water temp. with recent increase of HAB
Toxicity Analysis	<ul style="list-style-type: none"> ➤ Toxicity analysis of HAB using bioassay 	<ul style="list-style-type: none"> ➤ Toxicity analysis with high performance liquid chromatography and mass chromatography ➤ Effectiveness of ELISA method 	<ul style="list-style-type: none"> ➤ Toxicity analysis of <i>C. polykrikoides</i>, <i>Alexandrium</i> spp., <i>Microcystis</i> spp. and <i>Pseudo-nitzschia</i> spp. 	<ul style="list-style-type: none"> ➤ Toxicity analysis of different genetic populations of <i>A. tamarense</i>
Taxonomy	<ul style="list-style-type: none"> ➤ Identification of dinoflagellates by two dimension proteome reference map ➤ Molecular identification of different <i>Alexandrium</i> spp. strains 	<ul style="list-style-type: none"> ➤ Development of molecular biological approach to distinguish plankton population 	<ul style="list-style-type: none"> ➤ Ultrastructure and phylogeny of micro algae by molecular biological approach 	<ul style="list-style-type: none"> ➤ Identification of <i>A. tamarense</i> subpopulations by molecular biological approach
Mitigation measures	<ul style="list-style-type: none"> ➤ Coagulation rate of clay with HAB species ➤ Monitoring and forecasting of HAB by remote sensing ➤ Control of HAB using yellow clay and surfactant 	<ul style="list-style-type: none"> ➤ Biological control of HAB using viruses, bacteria, macroalgae ➤ HAB prediction with neural network technique 	<ul style="list-style-type: none"> ➤ Early detection of HAB with molecular biological technique ➤ Biological control of HAB using bacteria, parasites, copepods and ciliates ➤ Control of HAB using yellow clay and surfactant ➤ Environmental impact of control agents ➤ Red tide detection using satellite remote sensing 	Information N/A

5. Training Activity to Cope with HAB

5.1 Training Activities in the NOWPAP Region

Table 10 shows types of training courses conducted by the NOWPAP Members. Majority of the training courses were related to red tides, shellfish monitoring and HAB mitigation with main participants from monitoring organizations, research institutes and universities. China operates some training courses for different trainee groups. Japan has invited local fishermen and aquaculture operators into these training programs, since their participation is vital for HAB monitoring and mitigation. Korea has conducted red tide training courses for technicians of developing countries, together with various other training courses. There are no national HAB training programs in Russia yet.

Table 10 Types of HAB Training Courses Conducted in the NAWPAP Region

	Targeted personnel	Host organization	Subject
China	Personnel involved in red tide and shellfish poison in the monitoring centers of SOA	Information N/A	Lectures on red tide monitoring and toxin analysis
	Personnel from universities and research institutes involved in red tide and shellfish poison research	Information N/A	Lectures on HPLC technique for PSP and DSP detection
	Personnel from universities and research institutes involved in fisheries research	Information N/A	Lectures and discussions on disease control in aquaculture farms
	Personnel from coastal local governments involved in environmental monitoring including red tide	Information N/A	Lectures on red tide monitoring, species identification and toxin analysis
Japan	Technicians of local government fisheries laboratory	Japan Fisheries Resource Conservation Association (JFRCA)	Lectures on latest HAB information. Exercises in sampling, sample preservation, species identification, toxin analysis etc.
	Local fishermen and aquaculture operators	Local Governments	Lectures on HAB mechanism, mitigation measures, monitoring etc.
Korea	Technicians of developing countries	Korean International Cooperation Agency (KOICA)	Lectures on red tide monitoring and mitigation.
	Personnel involved in coastal zone management in local government or regional maritime affairs & fisheries	NFRDI	Lectures on red tide monitoring and mitigation. Lectures on HAB mechanism
	Technicians involved in sanitation and inspection of fishery products. Personnel from private fishery companies	NFRDI	Lectures on shellfish poison
	Personnel involved in red tide monitoring in regional maritime affairs & fisheries	NFRDI	Exercises in sampling, sample preservation, species identification, toxin analysis etc.

5.2 Common Issues on Training Activities in the NOWPAP Region

Each NOWPAP Member has realized the importance of capacity building for improving HAB monitoring (Report of the First Meeting of NOWPAP Working Group 3), and China, Japan and Korea have conducted various training programs; however, these countries have carried out the promotion of concrete techniques for HAB monitoring practice and toxin analysis in their own training programs. Since the NOWPAP Members conduct these training programs individually, there could be some differences in knowledge and techniques of the trainees. Therefore, to share common knowledge, to unite techniques for HAB monitoring and toxin analysis, and to implement a common HAB monitoring training program are an option for NOWPAP.

6. Suggested Activities for HAB in the NOWPAP Region

6.1 National Activities to Cope with HAB

According to National Report on HAB of each country, they conduct one or more national activities concerning HAB. Table 11 shows national activities that are currently implemented to cope with HAB problems.

Table 11 Implemented National Activities to Cope with HAB in the NOWPAP Region

China	Japan	Korea	Russia
<ul style="list-style-type: none"> ➤ Regular monitoring of red tide ➤ Use of clay spraying to control HAB (only in limited areas) 	<ul style="list-style-type: none"> ➤ Regular monitoring of red tide, toxin-producing plankton and shellfish poison ➤ Operation of HAB database (includes information on past HAB events) ➤ Use of clay spraying to control HAB (only in limited areas) 	<ul style="list-style-type: none"> ➤ Regular monitoring of red tide and shellfish poison. ➤ Dissemination of HAB information to concerned organizations and fishermen through the HAB Emergency Center ➤ Use of clay dispenser and electric clay dispenser to control <i>Cochlodinium</i> blooms ➤ Use of automatic HAB alarm system in aquaculture farms for early detection of HAB 	<ul style="list-style-type: none"> ➤ No national programs implemented yet apart from HAB related research activities.

Monitoring of red tide is currently implemented in Japan, China and Korea. Korea has a well established national monitoring scheme through the NFRDI, whereas in Japan and China, monitoring is not conducted under a national scheme.

Clay spraying is a common red tide mitigation method employed in China, Japan and Korea. Its use is limited to certain areas and situations because being concerned with negative effects to the environment. Korea has developed an automatic HAB alarm system, which provides early red tide warnings to fishermen.

Slight differences in the proposed national activities are inevitable among the NOWPAP Members since each NOWPAP Member has their own particular problems and priorities for HAB. For example, Russia's HAB monitoring system is still in its development stage, in which administrative reform is a priority for future development. On the other hand, Korea already has a well developed HAB monitoring system based on the Integrated Coastal Zone Management

Strategy.

In principal, all the NOWPAP Members have their own priorities on developing a more effective monitoring system and mitigation measure. The use of satellite remote sensing is considered as an effective tool for red tide monitoring by all the NOWPAP Members, and many research activities have been focused on this area. Biological control of HAB is another option studied by some of the NOWPAP Members.

6.2 Suggested Activity for HAB in the NOWPAP Region

National Report on HAB of each country made suggestions for NOWPAP future activities concerning HAB. Table 12 lists suggested future activities for HAB in the NOWPAP Region.

Table 12 Suggested Future Activities for HAB in the NOWPAP Region

China	Japan	Korea	Russia
<ul style="list-style-type: none"> ➤ Development of common data and information network for HAB monitoring (C1) ➤ Cooperation and exchange of information with other relevant organizations such as WESTPAC and PICES (C2) 	<ul style="list-style-type: none"> ➤ Action against <i>Cochlodinium</i> blooms, through continuation of CCG and organization of joint programs with WESTPAC/TTR (J1) ➤ Cooperation with other UNEP Action Plans (e.g. East Asia Sea Action Plan) (J2) ➤ Exchange of information with other organizations to avoid unnecessary overlaps of activities (J3) ➤ Development of appropriate policies and technologies to control inputs of land-based pollutants into the seas of the NOWPAP Region (J4) 	<ul style="list-style-type: none"> ➤ Action against <i>Cochlodinium</i> blooms, through continuation of CCG and implementation of collaborative research programs within the NOWPAP Members (K1) ➤ Development of appropriate policies and technologies to control inputs of land-based pollutants into the seas of the NOWPAP Region (K2) 	<ul style="list-style-type: none"> ➤ Research and analysis on the influence of land-based nutrients and pollutants on HABs in coastal zone. (R1) ➤ Cooperation and exchange of information with other relevant organizations such as WESTPAC and PICES (R2) ➤ Continuation of international training programs (R3)

Japan and Korea consider activities for *Cochlodinium* to be important. Damages of this species are severe on fisheries in these countries. Area of *Cochlodinium* occurrence tends to grow bigger in the NOWPAP Region. Even though the damages are not found in China and Russia in the NOWPAP Region for the time being, this species might be problems in the future. Therefore, the NOWPAP Member should treat *Cochlodinium* as a common problem for NOWPAP and cooperate with one another to conduct activities concerning this species. In 2004, *Cochlodinium* Corresponding Group (CCG) started a cooperative work on this species. This group activity should be continued and become more effective and cooperative. (Refer to Suggestion1 in Table 13)

China, Japan and Russia emphasize the importance of cooperation within the NOWPAP Region as well as other international organizations that are involved in HAB, such as IOC/WESTPAC and PICES. Valuable information could be exchanged, and activities could be demarcated through the process. Some objectives of this cooperation are to avoid overlapped activities done in each researcher through exchanging valuable information, and this wider range of information enables WG3 activities on HAB more active and leads them to solve HAB problems. (Refer to Suggestion2 in Table 13)

China suggests development of common data and information network for HAB monitoring. China has developed “China Harmful Algal Bloom WebPages (www.china-hab.cn)” and a website of the “National Basic Research Priority Project - China Ecology and Oceanology of Harmful Algae Blooms”. The latter website is embedded in the former one. These information systems are expected to enable prompt response to HAB occurrences and to accumulate scientific knowledge about HAB. Japan has constructed “Marine Environmental Watch Project” and “Website of Remote Sensing of the Japanese Coastal Guard” which provide satellite remote sensing images of chlorophyll-a. These data can be useful to investigate HABs. NOWPAP WG3 has developed “HAB Reference Database” and “Cochlodinium Homepage”. The former provides scientific reference information about HAB to the NOWPAP Members, and the latter introduces *Cochlodinium* which is one of the most concerned HAB species in the NOWPAP Region. Through more development of such a database and information network for NOWPAP, common understanding of HAB should be deepened. (Refer to Suggestion3 in Table 13)

Japan, Korea and Russia think that more effective policies and technologies are needed to control the discharge of land-based nutrients (e.g. effluent control and improvement of sewage treatment system). In order to help policy makers to implement new policies and encourage private sect to invent new technologies, NOWPAP WG3 may provide data of nutrient sources, river water quality, or nutrient loads in cooperation with NOWPAP WG1 and WG2, and give information about preventive measures which the NOWPAP Members have conducted since 1970s. (Refer to Suggestion4 in Table 13)

It is desirable that a collaborative monitoring program is developed within the NOWPAP Region to construct a common knowledge of HAB in this region. In reality, however, each country has already established their own programs with a long experience, which is difficult to change their own way; and they have used different definitions of words (e.g. name of species). It is really challenging to develop a collaborative monitoring program in this region but NOWPAP WG3 should make efforts to construct quasi-collaborative monitoring program with

feasible activities to share common information about HAB among the NOWPAP Members. This is not mentioned in National Reports but this is an ultimate goal of NOWPAP WG3. (Refer to Suggestion5 in Table 13)

Suggestions for future activities about HAB in the NOWPAP Region are summarized in Table 13. Five suggestions are made for WG3 future activities.

Table 13 Summary of the suggestions for future activities about HAB in the NOWPAP Region

- 1. To facilitate research and study of *Cochlodinium* through CCG activities (J1,K1)**
- 2. To cooperate with other international organizations that are involved in HAB (C2,J2,J3, R2, R3)**
- 3. To establish common understanding of HAB through development of a database and information network (C1)**
- 4. To help make a policy on control of land-based nutrients discharge (J4, K2, R1)**
- 5. To seek a collaborative work for HAB monitoring for the NOWPAP Region**

Considering the priority for WG3 for the next years among the five suggestions above, an action for Suggestion 1 and 4 come on top as “promotion of mitigation.” Until having developed knowledge of HAB in the NOWPAP Region through working on National Reports, Integrated Report, HAB Reference Database and CCG activities in the past four years, WG3 could not start activities for “promotion of mitigation” which was listed on the workplan proposed in FPMs and WG3 Meetings as an important action to be taken. Therefore, now may be a right time to take an action on “promotion of mitigation.” Also, collecting case studies of mitigation measures might be an option for WG3 activities for the following years.

The immediate topic for future cooperative work of CCG is to establish countermeasures against damages by *Cochlodinium* red tides. The present report describes current mitigation methods to prevent damages caused by *Cochlodinium* red tides. It should be noted that these mitigations have a very limited effect, and red tide cases are still increasing. It means that further development of countermeasures is necessary for sound development of utilization of the NOWPAP coastal region and conservation of its environment.

The need to establish effective countermeasures against HAB is not only limited to *Cochlodinium*, but is also to other HAB species as well. Some NOWPAP Members have already been implementing some mitigation measures against HAB, though with varying efforts and methods. Researches on this field are also an ongoing process by the NOWPAP Members.

All the NOWPAP Members must have preventive measures to mitigate Red-tide occurrence such as control of nutrient discharges. Especially Japan has a long experience of nutrient control since 1970 to implement laws and set standard on water qualities of effluents, rivers and sea areas. It is important to share information of preventive measures conducted in each of the NOWPAP Members in order to make better policies on control of land-based nutrient discharges.

In summary, considering the importance of HAB mitigation, one of the primary future activities of WG3 may be collection and compilation of detailed information related to HAB mitigation measures. This information includes both preventive measures (e.g. water and sediment quality standards, laws and regulation, etc.) and countermeasures (e.g. clay spraying) against red tide or HAB.

Appendices

- i Occurrences
of red tides in the NOWPAP Region (under construction)
- ii Red tide events in the NOWPAP Region (under construction)




Class	Genus and Species	Abbreviation
Cyanophyceae	<i>Microcystis viridis</i>	M.v.
Cryptophyceae	<i>Chromonas marina</i>	Chro.m.
	<i>Chromonas salina</i>	Chro.s.
	<i>Cryptomonas acuta</i>	Cry.a.
Dinophyceae	<i>Cryptomonas</i> sp.	Cry.sp.
	<i>Alexandrium catenella</i>	A.c.
	<i>Alexandrium fraterculus</i>	A.f.
	<i>Alexandrium</i> sp.	A.sp.
	<i>Ceratium furca</i>	C.f.
	<i>Ceratium fusus</i>	C.fusus
	<i>Ceratium</i> sp.	C.sp.
	<i>Cochlodinium polykrikoides</i>	C.p.
	<i>Cochlodinium</i> sp.	Coch.sp.
	<i>Gymnodinium mikimotoi</i>	G.m.
	<i>Gymnodinium sanguineum</i>	G.s.
	<i>Gymnodinium</i> sp.	Gym.sp.
	<i>Gyrodinium</i> sp.	Gvro.sp.
	<i>Heterocapsa circularisquama</i>	H.c.
	<i>Heterocapsa triquetra</i>	H.t.
	<i>Heterocapsa</i> sp.	Heteroc.sp.
	<i>Noctiluca scintillans</i>	N.s.
	<i>Noctiluca</i> sp.	Noc.sp.
	<i>Oxyrrhis marina</i>	O.m.
	<i>Prorocentrum balticum</i>	P.b.
	<i>Prorocentrum dentatum</i>	P.d.
<i>Prorocentrum micans</i>	P.mic.	
<i>Prorocentrum minimum</i>	P.m.	
<i>Prorocentrum sigmoides</i>	P.s.	
<i>Prorocentrum triestinum</i>	P.t.	
<i>Prorocentrum</i> sp.	Pro.sp.	
Chrysophyceae	<i>Dictyocha fibula</i>	D.f.
Haptophyceae	<i>Phaeocystis</i> sp.	Phaeo.sp.
Haptophyceae	<i>Haptophyceae</i>	Hapto.
Bacillariophyceae	<i>Asterionella</i> sp.	Aste.sp.
	<i>Chaetoceros curvisetum</i>	Ch.c.
	<i>Chaetoceros pseudocurvisetum</i>	Ch.ps.
	<i>Chaetoceros socialis</i>	Ch.s.
	<i>Chaetoceros</i> sp.	Chaeto.sp.
	<i>Coscinodiscus gigas</i>	Cos.g.
	<i>Coscinodiscus</i> sp.	Cos.sp.
	<i>Ditylum brightwellii</i>	Dity.b.
	<i>Eucampia zodiacus</i>	Euc.z.
	<i>Leptocylindrus danicus</i>	L.d.
	<i>Leptocylindrus</i> sp.	Lepto.sp.
	<i>Navicula</i> sp.	Nav.sp.
	<i>Neodelphineis pelagica</i>	Neo.p.
	<i>Nitzschia pungens</i>	N.p.
	<i>Nitzschia</i> sp.	Nit.sp.
	<i>Nitzschia</i> spp.	Nitz.spp.
	<i>Pseudo-nitzschia pungens</i>	Pse.p.
	<i>Pseudo-nitzschia</i> sp.	Pse.sp.
	<i>Rhizosolenia delicatula</i>	R.d.
	<i>Rhizosolenia fragilissima</i>	R.f.
	<i>Rhizosolenia</i> sp.	Rhizo.sp.
<i>Skeletonema costatum</i>	S.c.	
<i>Skeletonema</i> sp.	S.sp.	
<i>Thalassiosira decipiens</i>	T.d.	
<i>Thalassiosira rotula</i>	T.r.	
<i>Thalassiosira</i> sp.	Thala.sp.	
Raphidophyceae	<i>Chattonella antiqua</i>	C.a.
	<i>Chattonella globosa</i>	C.g.
	<i>Chattonella marina</i>	C.m.
	<i>Fibrocapsa japonica</i>	F.j.
	<i>Heterosigma akashiwo</i>	H.a.
	<i>Heterosigma</i> sp.	H.sp.
Euglenophyceae	<i>Eutreptia lanowii</i>	E.l.
	<i>Eutreptiella</i> sp.	E.sp.
Prasinophyceae	<i>Pyramimonas</i> sp.	Pyra.sp.
Ciliate	<i>Mesodinium rubrum</i>	M.r.
	<i>Tontonia</i> sp.	Ton.sp.

Location of red tides in the NOWPAP Region (1)



Class	Genus and Species	Abbreviation
Cyanophyceae	<i>Microcystis viridis</i>	M.v.
Cryptophyceae	<i>Chromonas marina</i>	Chro.m.
	<i>Chromonas salina</i>	Chro.s.
	<i>Cryptomonas acuta</i>	Cry.a.
	<i>Cryptomonas</i> sp.	Cry.sp.
Dinophyceae	<i>Alexandrium catenella</i>	A.c.
	<i>Alexandrium fraterculus</i>	A.f.
	<i>Alexandrium</i> sp.	A.sp.
	<i>Ceratium furca</i>	C.f.
	<i>Ceratium fuscus</i>	C.fuscus
	<i>Ceratium</i> sp.	C.sp.
	<i>Cochlodinium polykrikoides</i>	C.p.
	<i>Cochlodinium</i> sp.	Coch.sp.
	<i>Gymnodinium mikimotoi</i>	G.m.
	<i>Gymnodinium sansuineum</i>	G.s.
	<i>Gymnodinium</i> sp.	Gym.sp.
	<i>Gyrodinium</i> sp.	Gyro.sp.
	<i>Heterocapsa circularisquama</i>	H.c.
	<i>Heterocapsa triquetra</i>	H.t.
	<i>Heterocapsa</i> sp.	Heteroc.sp.
	<i>Noctiluca scintillans</i>	N.s.
	<i>Noctiluca</i> sp.	Noc.sp.
	<i>Oxyrrhis marina</i>	O.m.
	<i>Prorocentrum balticum</i>	P.b.
	<i>Prorocentrum dentatum</i>	P.d.
<i>Prorocentrum micans</i>	P.mic.	
<i>Prorocentrum minimum</i>	P.m.	
<i>Prorocentrum sigmoides</i>	P.s.	
<i>Prorocentrum trisetinum</i>	P.t.	
<i>Prorocentrum</i> sp.	Pro.sp.	
Chrysophyceae	<i>Dictyocha fibula</i>	D.f.
Haptophyceae	<i>Phaeocystis</i> sp.	Phaeo.sp.
	Haptophyceae	Hapto.
Bacillariophyceae	<i>Asterionella</i> sp.	Aste.sp.
	<i>Chaetoceros curvisetum</i>	Ch.c.
	<i>Chaetoceros pseudocurvisetum</i>	Ch.ps.
	<i>Chaetoceros socialis</i>	Ch.s.
	<i>Chaetoceros</i> sp.	Chaeto.sp.
	<i>Coscinodiscus gigas</i>	Cos.g.
	<i>Coscinodiscus</i> sp.	Cos.sp.
	<i>Dietylum brightwellii</i>	Diety.b.
	<i>Eucampia zodiacus</i>	Euc.z.
	<i>Leptocylindrus danicus</i>	L.d.
	<i>Leptocylindrus</i> sp.	Lepto.sp.
	<i>Navicula</i> sp.	Nav.sp.
	<i>Neodelphineis pelagica</i>	Neo.p.
	<i>Nitzschia pungens</i>	N.p.
	<i>Nitzschia</i> sp.	Nit.sp.
	<i>Nitzschia</i> spp.	Nitz.spp.
	<i>Pseudo-nitzschia pungens</i>	Pse.p.
	<i>Pseudo-nitzschia</i> sp.	Pse.sp.
	<i>Rhizosolenia delicatula</i>	R.d.
	<i>Rhizosolenia fragilissima</i>	R.f.
	<i>Rhizosolenia</i> sp.	Rhizo.sp.
	<i>Skeletonema costatum</i>	S.c.
	<i>Skeletonema</i> sp.	S.sp.
<i>Thalassiosira decipiens</i>	T.d.	
<i>Thalassiosira rotula</i>	T.r.	
<i>Thalassiosira</i> sp.	Thala.sp.	
Raphidophyceae	<i>Chattonella antiqua</i>	C.a.
	<i>Chattonella globosa</i>	C.g.
	<i>Chattonella marina</i>	C.m.
	<i>Fibrocapsa japonica</i>	F.j.
	<i>Heterosigma akashiwo</i>	H.a.
	<i>Heterosigma</i> sp.	H.sp.
Euglenophyceae	<i>Eutreptia lanowii</i>	E.l.
	<i>Eutreptiella gymnastica</i>	E.g.
	<i>Eutreptiella</i> sp.	E.sp.
Prasinophyceae	<i>Pyramimonas</i> sp.	Pyra.sp.
Ciliate	<i>Mesodinium rubrum</i>	M.r.
	<i>Tontonia</i> sp.	Ton.sp.

Location of red tides in the NOWPAP Region (2)



Class	Genus and Species	Abbreviation
Cyanophyceae	<i>Microcystis viridis</i>	M.v.
Cryptophyceae	<i>Chromonas marina</i>	Chro.m.
	<i>Chromonas salina</i>	Chro.s.
	<i>Cryptomonas acuta</i>	Cry.a.
	<i>Cryptomonas</i> sp.	Cry.sp.
Dinophyceae	<i>Alexandrium catenella</i>	A.c.
	<i>Alexandrium fraterculus</i>	A.f.
	<i>Alexandrium</i> sp.	A.sp.
	<i>Ceratium furca</i>	C.f.
	<i>Ceratium fuscus</i>	C.fuscus
	<i>Ceratium</i> sp.	C.sp.
	<i>Cochlodinium polykrikoides</i>	C.p.
	<i>Cochlodinium</i> sp.	Coch.sp.
	<i>Gymnodinium mikimotoi</i>	G.m.
	<i>Gymnodinium sanguineum</i>	G.s.
	<i>Gymnodinium</i> sp.	Gym.sp.
	<i>Gyrodinium</i> sp.	Gyro.sp.
	<i>Heterocapsa circularisquama</i>	H.c.
	<i>Heterocapsa triquetra</i>	H.t.
	<i>Heterocapsa</i> sp.	Heteroc.sp.
	<i>Noctiluca scintillans</i>	N.s.
	<i>Noctiluca</i> sp.	Noc.sp.
	<i>Oxyrrhis marina</i>	O.m.
	<i>Prorocentrum balticum</i>	P.b.
	<i>Prorocentrum dentatum</i>	P.d.
<i>Prorocentrum micans</i>	P.mic.	
<i>Prorocentrum minimum</i>	P.m.	
<i>Prorocentrum sigmoides</i>	P.s.	
<i>Prorocentrum triestinum</i>	P.t.	
<i>Prorocentrum</i> sp.	Pro.sp.	
Chrysophyceae	<i>Dictyocha fibula</i>	D.f.
Haptophyceae	<i>Phaeocystis</i> sp.	Phaeo.sp.
Haptophyceae	<i>Haptophyceae</i>	Hapto.
Bacillariophyceae	<i>Asterionella</i> sp.	Aste.sp.
	<i>Chaetoceros curvisetum</i>	Ch.c.
	<i>Chaetoceros pseudocurvisetum</i>	Ch.ps.
	<i>Chaetoceros socialis</i>	Ch.s.
	<i>Chaetoceros</i> sp.	Chaeto.sp.
	<i>Coscinodiscus gigas</i>	Cos.g.
	<i>Coscinodiscus</i> sp.	Cos.sp.
	<i>Ditylum brightwellii</i>	Dity.b.
	<i>Eucampia zodiacus</i>	Euc.z.
	<i>Leptocylindrus danicus</i>	L.d.
	<i>Leptocylindrus</i> sp.	Lepto.sp.
	<i>Navicula</i> sp.	Nav.sp.
	<i>Neodelphineis pelagica</i>	Neo.p.
	<i>Nitzschia pungens</i>	N.p.
	<i>Nitzschia</i> sp.	Nit.sp.
	<i>Nitzschia</i> spp.	Nitz.spp.
	<i>Pseudo-nitzschia pungens</i>	Pse.p.
	<i>Pseudo-nitzschia</i> sp.	Pse.sp.
	<i>Rhizosolenia delicatula</i>	R.d.
	<i>Rhizosolenia fragilissima</i>	R.f.
<i>Rhizosolenia</i> sp.	Rhizo.sp.	
<i>Skeletonema costatum</i>	S.c.	
<i>Skeletonema</i> sp.	S.sp.	
<i>Thalassiosira decipiens</i>	T.d.	
<i>Thalassiosira rotula</i>	T.r.	
<i>Thalassiosira</i> sp.	Thala.sp.	
Raphidophyceae	<i>Chattonella antiqua</i>	C.a.
	<i>Chattonella globosa</i>	C.g.
	<i>Chattonella marina</i>	C.m.
	<i>Fibrocapsa japonica</i>	F.j.
	<i>Heterosigma akashiwo</i>	H.a.
<i>Heterosigma</i> sp.	H.sp.	
Euglenophyceae	<i>Eutreptia lanowii</i>	E.l.
	<i>Eutreptiella gymnastica</i>	E.g.
Euglenophyceae	<i>Eutreptiella</i> sp.	E.sp.
Prasinophyceae	<i>Pyramimonas</i> sp.	Pyra.sp.
Ciliate	<i>Mesodinium rubrum</i>	M.r.
	<i>Tontonia</i> sp.	Ton.sp.

Location of red tides in the NOWPAP Region (3)

Red tide events in China (B Sea Area and C Sea Area) (1)

Event No.	Location (name of the sea area)	Duration dd/mm/yy-dd/mm/yy	Causative species	Max. cell density (cells/L)	Approximate Area suffered (km ²)	Type of HAB Red tide or Toxic	Mitigation Activity and effectiveness	Damage	
								Fishery resources	Human health
1	Huanghua, Hebei	1989	<i>Gymnodinium</i> sp	No data	1,300	HAB	No data	38 million dollar	No data
2	Laizhou Bay	18/6/1990	No data	No data	1/3 Bay area	Red tide	No data	No data	No data
3	Jiaozhou Bay	26/6/1990	No data	No data	80,000	Red tide	No data	No data	No data
4	Baidaihe, Hebei	28/6/1990-4/7/1990	No data	No data	110	No data	No data	No data	No data
5	Laizhou Bay	June, 1990	<i>Noctiluca scintillans</i>	No data	No data	Red tide	No data	No data	No data
6	Laizhou Bay	19-20/8/1990	No data	No data	10	Red tide	No data	No data	No data
7	Laizhou Bay	26/8/1990	No data	No data	1,200	Red tide	No data	No data	No data
8	Laizhou Bay	30/8/1990	No data	No data	1,000	Red tide	No data	No data	No data
9	North Laizhou Bay	1/9/1990	No data	No data	No data	Red tide	No data	No data	No data
10	Changhai county, Liaoning	1990	No data	No data	No data	HAB	No data	2.5 million dollar due to death scallops	No data
11	Shrimp pond, Dalian	May to July, 1991	<i>Exuviaella cordata</i>	7.5 10 ⁷	No data	Red tide	No data	Loss of shrimp	No data
12	Liaodong Bay	4/7/1991-12/7/1991	<i>Noctiluca scintillans</i>	4.9 10 ⁷	100	Red tide	No data	No data	No data
13	Jiaozhou Bay	April, 1992	No data	No data	No data	Red tide	No data	No data	No data
14	East Qingdao	12/5/1992	No data	No data	1,200	Red tide	No data	No data	No data
15	Jiaozhou Bay	August, 1992	No data	No data	1,000	Red tide	No data	No data	No data
16	Dalian Bay	11/8/1993	No data	No data	40	Red tide	No data	No data	No data
17	Laizhou Bay	6/6/1995	<i>Noctiluca scintillans</i>	2.16 10 ⁷	90	Red tide	No data	No data	No data
18	Liaodong Bay	20/8/1995	No data	No data	100	Red tide	No data	No data	No data
19	Penglai, Laizhou Bay	13-14/4/1997	No data	No data	1	Red tide	No data	No data	No data
20	Bohai Bay	28/6/1997	No data	No data	3	Red tide	No data	No data	No data
21	Jiaozhou Bay	3-8/7/1998	<i>Skeletonema costatum</i>	4.5 10 ⁶	10	Red tide	No data	No data	No data
22	Yantai, Laizhou Bay	August, 1998	No data	No data	100	HAB	No data	4 million dollar Fishery losses	
23	Bohai Sea	16/8/1998-19/9/1998	<i>Ceratium furca</i> , <i>Dinophysis ovata</i>	1.25 10 ⁶	5,000	toxic	No data	15 million dollar Fishery losses	DSP detected
24	Yantai, Bohai	15/8/1998-10/9/1998	<i>Gymnodinium sanguineum</i>	No data	170	HAB	No data	Shellfish death	No data
25	Laizhou Bay	2/9/1998	No data	No data	No data	Red tide	No data	No data	No data
26	Liaodong Bay	18/9/1998	<i>Ceratium furca</i>	No data	No data	Red tide	No data	No data	No data
27	Liaodong Bay	29/9/1998	<i>Ceratium furca</i>	No data	No data	Red tide	No data	No data	No data
28	Bohai Bay	1/10/1998	No data	No data	No data	Red tide	No data	No data	No data
29	Bohai Bay	3/10/1998	<i>Gonyaulax spinifera.</i> , <i>Ceratium furca</i>	No data	800	Red tide	No data	No data	No data
30	Bohai Bay	9/10/1998	No data	No data	No data	Red tide	No data	No data	No data
31	Jiaozhou Bay	8-15/6/1999	<i>Eucampia zoodiacus</i>	2.3 10 ⁶	No Data	Red tide	No data	No data	No data
32	Bohai Bay	2-4/7/1999	No data	No data	1,500	Red tide	No data	No data	No data
33	Dalian Bay	July, 1999	<i>Exuviaella marina</i>	8.1 10 ⁶	No data	HAB	No data	No data	DSP detected
34	Bohai Sea	13-21/7/1999	<i>Noctiluca scintillans</i>	No data	6,300	Red tide	No data	No data	No data
35	Dalian Bay	17-21/7/1999	<i>Noctiluca scintillans</i>	No data	100	Red tide	No data	No data	No data
36	Penglai, Shandong	17/7/1999	<i>Noctiluca scintillans</i>	No data	680	Red tide	No data	No data	No data
37	South Dalian	18/7/1999	No data	No data	30	Red tide	No data	No data	No data
38	Jiaozhou Bay	23/7/1999	<i>Skeletonema costatum</i>	No data	26	Red tide	No data	No data	No data

Red tide events in China (B Sea Area and C Sea Area) (2)

Event No.	Location (name of the sea area)	Duration dd/mm/yy-dd/mm/yy	Causative species	Max. cell density (cells/L)	Approximate Area suffered (km ²)	Type of HAB Red tide or Toxic	Mitigation Activity and effectiveness	Damage	
								Fishery resources	Human health
39	Xiaomai Island, Qingdao	26/7/1999	No data	No data	60	Red tide	No data	No data	No data
40	Shidao, Shangdong	6/8/1999	No data	No data	160	Red tide	No data	No data	No data
41	Central Bohai Sea	25/9/1999	No data	No data	30	Red tide	No data	No data	No data
42	Liaodong Bay, Bohai	9-15/7/2000	<i>Noctiluca scintillans</i>	No data	350	Red tide	No data	No data	No data
43	Liaodong Bay	Jul-00	<i>Prorocentrum</i> sp.	No data	No data	HAB	No data	Death of jellyfish	No data
44	Bohai Bay	23/7/2000	No data	No data	1,040	Red tide	No data	No data	No data
45	North Wentuozhi Island, Bohai	13/8/2000	No data	No data	217	Red tide	No data	No data	No data
46	Changxin Island, Bohai Sea	13/8/1/2000	No data	No data	44	Red tide	No data	No data	No data
47	Zhuanghe, Yellow Sea	2/8/2000	No data	No data	827	HAB	No data	15 million	No data
48	Southeast Qikou	20-21/7/2000	No data	No data	180	Red tide	No data	No data	No data
49	Beidaihe, Tianjing	23/7/2000	No data	No data	3	Red tide	No data	No data	No data
50	Tangu, Tianjing	25/7/2000	No data	No data	134	Red tide	No data	No data	No data
51	Jiaozhou Bay	20-23/7/2000	<i>Noctiluca scintillans</i>	No data	2	Red tide	No data	No data	No data
52	Dandong, North Yellow Sea	24/5/2001	No data	No data	No data	Red tide	No data	No data	No data
53	Bohai Bay	26/5/2001	No data	No data	No data	Red tide	No data	No data	No data
54	Bohai Bay	19/6/2001	No data	No data	No data	Red tide	No data	No data	No data
55	Jiaozhou Bay	11-12/6/2001	<i>Noctiluca scintillans</i>	No data	5	Red tide	No data	No data	No data
56	The Coast of Jiangsu	20/6/2001	<i>Skeletonema costatum</i>	No data	1,000	Red tide	No data	No data	No data
57	Jiaozhou Bay	7-12/7/2001	<i>Mesodinium rubrum</i>	No data	20	Red tide	No data	No data	No data
58	Yingkou, Liaodong	15-16/7/2001	<i>Noctiluca scintillans</i>	No data	360	Red tide	No data	No data	No data
59	Bayuquan, Liaodong Bay	12-23/8/2001	<i>Leptocylindrus danicus</i>	No data	770	Red tide	No data	No data	No data
60	Yalujiang Estuary, North Yellow Sea	24/8/2001-14/9/2001	<i>Eucampia zoodiacus, Chaetocerus socialis</i>	No data	1,100	Red tide	No data	No data	No data
61	Liao River Estuary	25-26/8/2001	<i>Navicula</i> sp.	No data	130	Red tide	No data	No data	No data
62	Bayuquan, Liaodong Bay	27-30/8/2001	<i>Mesodinium rubrum, Eucampia zoodiacus</i>	No data	100	Red tide	No data	No data	No data
63	Qinghuangdao Bay, Bohai Sea	3-4/6/2002	<i>Noctiluca scintillans</i>	No data	1	Red tide	No data	No data	No data
64	Jingtang Harbour, Bohai Bay	16-17/6/2002	<i>Noctiluca scintillans</i>	No data	15	Red tide	No data	No data	No data
65	Jingtang Harbour, Bohai Bay	27/6/2002	<i>Gymnodinium</i> sp., <i>Noctiluca scintillans</i> ,	No data	1	Red tide	No data	No data	No data
66	Qinghuangdao Bay, Bohai Sea	25/7/2002	<i>Chattonella marina</i>	No data	8	HAB	No data	No data	No data
67	Laizhou Bay	10/8/2002	<i>Noctiluca scintillans</i>	No data	20	HAB	No data	0.6 million	No data
68	Laizhou Bay	15/8/2002	<i>Skeletonema costatum</i>	No data	30	HAB	No data	1 million dollar	No data
69	East Liaodong Bay	28/5/2003	<i>Noctiluca scintillans</i>	No data	10	Red tide	No data	No data	No data
70	Dandong waters, Yellow Sea	Jun-03	No data	No data	30	Red tide	No data	No data	No data

Red tide events in China (B Sea Area and C Sea Area) (3)

Event No.	Location (name of the sea area)	Duration dd/mm/yy-dd/mm/yy	Causative species	Max. cell density (cells/L)	Approximate Area suffered (km ²)	Type of HAB Red tide or Toxic	Mitigation Activity and effectiveness	Damage	
								Fishery resources	Human health
71	Dalian Bay	Jul-03	<i>Heterosigma akashiwo</i>	No data	15	HAB	No data	No data	No data
72	Jiaozhou Bay	Jul-03	<i>Coscinodiscus asteromphalus</i>	No data	200	Red tide	No data	No data	No data
73	Qinghuangdao, East Bohai Sea	25-26/4/2003-	<i>Noctiluca scintillans</i>	No data	70	Red tide	No data	No data	No data
74	Liaodong Bay	28/5/2003	<i>Noctiluca scintillans</i>	No data	10	Red tide	No data	No data	No data
75	Qinghuangdao, East Bohai Sea	28/5/2003-4/6/2003	<i>Noctiluca scintillans</i>	No data	8	Red tide	No data	No data	No data
76	Liaodong Bay	28/5/2003	<i>Noctiluca scintillans</i>	No data	140	HAB	No data	Fish kills	No data
77	Qinghuangdao, East Bohai Sea	12/6/2003	<i>Noctiluca scintillans</i>	No data	0	Red tide	No data	No data	No data
78	Luanhe, Qinghuangdao, East Bohai Sea	21/6/2003	<i>Noctiluca scintillans</i>	No data	12	Red tide	No data	No data	No data
79	Qinghuangdao, East Bohai Sea	25-27/6/2003	<i>Noctiluca scintillans</i>	No data	1	Red tide	No data	No data	No data
80	Dagu Harbour,	1-8/7/2003	<i>Noctiluca scintillans</i>	No data	100	Red tide	No data	No data	No data
81	Bohai Bay	12-13/8/2003	<i>Noctiluca scintillans</i>	No data	2	Red tide	No data	No data	No data
82	Laizhou Bay	2003	<i>Gonyaulax spinifera</i>	No data	No data	Red tide	No data	No data	No data
83	Jiaozhou Bay	9-28/2/2004	<i>Rhizosolenia</i> sp.	No data	No data	Red tide	No data	No data	No data
84	Yellow River Estuary	11-18/6/2004	<i>Phaeocystis</i> sp.	No data	1,850	HAB	No data	No data	No data
85	Central Bohai Bay	12-18/6/2004	<i>Kerania mikimotoi</i>	No data	3,200	HAB	No data	No data	No data
86	Jingshitan , Dalian , Yellow Sea	6/9/2004	<i>Chattonella antiqua</i>	No data	No data	HAB	No data	No data	No data
87	Jingshitan , Dalian , Yellow Sea	25/9/2004	<i>Alexandrium catenella</i>	No data	No data	HAB	No data	No data	No data

Red tide events in Japan (northern Kyushu coastal waters) (3)

Event No.	Location (name of the sea area)		Duration dd/mm/yy-dd/mm/yy	Approximate Area suffered (km ² -km)	Fish/shellfish species				Contents				Quantity				Economic loss(thousand yen)				Human health	
	Location 1	Location 2																				
NS-01	remote Island	Tsushima	27/01/98 - 02/02/98	0.75																		
NS-02	remote Island	Goto	07/04/98 - 25/04/98	0.5																		
NS-08	remote Island	Tsushima	16/05/98 - 21/05/98	0.002																		
FO-03	N	other	01/06/98 - 05/06/98	5																		
YG-01	N	other	03/06/98 - 16/06/98	0.06	Jackmackerel, Amberjacks					died				unknown								20 ~ 30
FO-04	N	Fukuokawan	15/06/98 - 17/06/98	71.1																		
SA-04	N	other	22/06/98 - 03/07/98	unknown																		
SA-05	N	other	24/06/98 - 26/06/98	unknown																		
SA-06	N	Imariwan	24/06/98 - 29/06/98	unknown																		
FO-06	N	Fukuokawan	13/07/98 - 16/07/98	41.5																		
YG-02	N	other	11/08/98 - 20/08/98	0.65	Fishes					died				unknown								unknown
NS-16	remote Island	Tsushima	17/08/98 - 21/08/98	1	Amberjacks					died				340 kg								1,122
YG-03	N	other	19/08/98 - 02/09/98	0.65																		
NS-17	remote Island	Tsushima	25/08/98 - 26/08/98	0.015																		
FO-08	N	Fukuokawan	27/08/98 - 28/08/98	40.6																		
FO-09	N	Fukuokawan	09/09/98 - 10/09/98	49																		
NS-22	N	Imariwan	31/10/98 - 24/10/98	3																		
NS-98-29	remote Island	Goto	17/12/98 - 04/01/99	0.995																		
NS-98-30	remote Island	Tsushima	28/12/98 - 06/01/99	0.035																		
NS-01	remote Island	Goto	05/01/99 - 09/01/99	0.285																		
YG-01	N	other	10/03/99 - 12/03/99	-	Squids, Octopus, fishes					died				13 kg								30
SA-01	N	other	04/04/99 - 20/04/99	-																		
NS-02	remote Island	Tsushima	19/04/99 - 26/04/99	2.51																		
YG-02	N	other	20/04/99 - 21/04/99	-																		
YG-03	N	other	20/04/99 - 21/04/99	0.992																		
YG-04	N	other	26/04/99 - 27/04/99	-																		
FO-02	N	Fukuokawan	10/05/99 - 12/05/99	75																		
YG-05	N	other	12/05/99 - 14/05/99	-																		
FO-03	N	Fukuokawan	31/05/99 - 02/06/99	30																		
SA-02	N	other	07/06/99 - 05/07/99	-																		
FO-04	N	other	08/06/99 - 10/06/99	1																		
FO-05	N	Fukuokawan	09/06/99 - 14/06/99	35																		
SA-03	N	other	20/06/99 - 26/07/99	-																		
YG-06	N	other	21/06/99 - 22/06/99	-																		
NS-09	N	Imariwan	01/07/99 - 21/07/99	4																		
FO-09	N	Fukuokawan	05/07/99 - 08/07/99	47																		
SA-04	N	Imariwan	05/07/99 - 29/07/99	-																		
SA-06	N	other	22/07/99 - 30/07/99	-																		
FO-11	N	Fukuokawan	22/07/99 - 22/08/99	62																		
NS-11	N	Imariwan	25/07/99 - 06/08/99	6																		
SA-07	N	Imariwan	03/08/99 - 09/08/99	-																		
SA-08	N	other	03/08/99 - 09/08/99	-																		
SA-09	N	other	05/08/99 - 09/08/99	-																		
NS-13	N	Imariwan	07/08/99 - 12/08/99	5	Sea bream	Yellowtail	Puffy fish	Horse mackerel		died	died	died	died	360,000 inds.	190,000 inds.	150,000 inds.	30,000 inds.	30,000 inds.	340,000	220,000	180,000	20,000
FO-13	N	other	09/08/99 - 18/08/99	1	Abalone					died				5,100 inds.								74
SA-10	N	Imariwan	10/08/99 - 16/08/99	15																		
SA-12	N	Imariwan	16/08/99 - 27/09/99	-																		
NS-24	remote Island	Tsushima	06/09/99 - 17/09/99	0.4																		
FO-16	N	Fukuokawan	07/09/99 - 13/09/99	40																		
NS-32	remote Island	Tsushima	09/12/99 - 21/12/99	0.07																		
YG-01	N	other	05/04/00	0.1																		
FO-02	N	other	23/05/00 - 26/05/00	under 1	Sea bream, Jackmackerel					died				unknown								unknown
FO-04	N	other	01/06/00 - 06/06/00	unknown																		
FO-05	N	Fukuokawan	02/06/00 - 06/06/00	65																		
FO-06	N	Fukuokawan	13/06/00 - 19/06/00	60																		
SA-03	N	other	15/06/00 - 19/06/00	0.005																		
SA-04	N	other	18/06/00 - 26/06/00	0.3	Amberjacks					died				400 inds.								400
SA-05	N	other	26/06/00 - 30/06/00	0.125																		
SA-06	N	Imariwan	27/06/00 - 27/07/00	4																		
FO-07	N	Fukuokawan	30/06/00 - 31/07/00	40																		
YG-02	N	other	06/07/00 - 03/08/00	unknown	Fishes, Abalone	Turban	Abalone	Turban	Turban	died	died	died	died	unknown	30 inds.	800kg	2000kg	unknown				
SA-07	N	other	10/07/00 - 19/07/00	2																		
NS-09	N	other	11/07/00 - 12/07/00	0.01																		
FO-09	N	other	11/07/00 - 31/07/00	unknown	Abalone, Turban					died				unknown								unknown
NS-10	N	Imariwan	13/07/00 - 22/07/00	16	Puffy fish					died				8,000 inds.								1,600
NS-12	remote Island	Goto	26/07/00 - 17/08/00	2	Turban					died				400 kg	120 inds.							340
FO-10	N	Fukuokawan	04/08/00 - 11/08/00	70																		
SA-10	N	Imariwan	18/08/00 - 11/09/00	unknown																		
NS-19	remote Island	Tsushima	21/08/00 - 24/08/00	0.2																		
FO-13	N	Fukuokawan	23/08/00 - 04/09/00	35																		
FO-14	N	Fukuokawan	08/09/00 - 12/09/00	unknown																		
NS-21	remote Island	Tsushima	18/09/00 - 25/09/00	0.0005																		
SA-11	N	Imariwan	27/09/00 - 29/09/00	12																		
NS-23	N	Imariwan	28/09/00 - 04/10/00	3																		
FO-15	N	Fukuokawan	07/11/00 - 11/11/00	unknown																		

Source : Kyushu Fishery Coordination Office, "Situation of Red Tide in the Seas surrounding Kyushu Island, 1999-2003.

Red tide events in Japan (northern Kyushu coastal waters) (4)

Event No.	Location (name of the sea area)		Duration dd/mm/yy-dd/mm/yy	Approximate Area suffered (km ² -km)	Fish/shellfish species				Contents			Quantity			Economic loss(thousand yen)			Human health
	Location 1	Location 2																
NS-27	remote Island	Tsushima	08/11/00 - 15/11/00	14														
SA-01	N	Imariwan	21/01/01 - 25/01/01	unknown														
YG-01	N	other	20/03/01 - 23/04/01	unknown														
FO-01	N	other	21/03/01 - 22/03/01	0.4														
FO-03	N	other	06/04/01 - 09/04/01	4														
NS-03	remote Island	Goto	07/04/01 - 11/04/01	2														
NS-04	N	Imariwan	10/04/01 - 13/04/01	0.1														
NS-05	remote Island	Iki	17/04/01 - 20/04/01	0.3														
NS-06	remote Island	Goto	17/04/01 - 18/04/01	58.7														
FO-04	N	other	17/04/01 - 20/04/01	60														
NS-07	remote Island	Tsushima	18/04/01 - 19/04/01	0.0025														
NS-08	remote Island	Tsushima	18/04/01 - 19/04/01	0.01														
SA-02	N	other	18/04/01 - 12/05/01	0.08														
NS-09	remote Island	Iki	27/04/01 - 01/05/01	unknown														
FO-05	N	Fukuokawan	06/05/01 - 14/05/01	70														
SA-03	N	other	07/05/01 - 11/05/01	0.4														
NS-12	remote Island	Goto	22/05/01 - 23/05/01	unknown														
YG-02	N	other	28/05/01 - 31/05/01	unknown														
NS-14	N	Imariwan	30/05/01 - 31/05/01	0.2														
FO-09	N	other	05/06/01 - 11/06/01	0.001														
YG-03	N	other	15/06/01	unknown														
NS-19	N	Imariwan	20/06/01 - 26/06/01	0.4														
FO-10	N	Fukuokawan	26/06/01 - 06/07/01	70														
YG-04	N	other	27/06/01 - 10/07/01	unknown														
NS-21	N	Imariwan	28/06/01 - 08/07/01	0.1														
FO-12	N	Fukuokawan	09/07/01 - 23/07/01	80														
NS-25	remote Island	Goto	19/07/01 - 24/07/01	unknown														
YG-05	N	other	03/08/01	unknown														
YG-06	N	other	06/08/01	0.02														
NS-32	remote Island	Tsushima	06/09/01 - 07/09/01	0.12														
FO-15	N	Fukuokawan	03/10/01 - 11/10/01	約10	Puffly fish	Amberjacks	Yellowtail	Fishes	died	died	died	226 kg	6 kg	3 kg		230	7	9
FO-16	N	other	01/11/01	under 1														
NS-40	remote Island	Tsushima	19/11/01 - 23/11/01	unknown														
FO-18	N	other	21/11/01 - 22/11/01	under 1														
NS-41	remote Island	Goto	03/12/01 - 05/12/01	0.15														
NS-42	remote Island	Tsushima	10/12/01	0.07														
NS-02	N	Imariwan	14/01/02 - 17/01/02	0.5														
YG-01	N	other	13/03/02 - 22/04/02	185														
FO-02	N	other	14/03/02	4														
NS-04	remote Island	Goto	01/04/02 - 02/04/02	2.4														
NS-06	remote Island	Goto	23/04/02	unknown														
NS-07	remote Island	Iki	24/04/02 - 26/04/02	unknown														
NS-10	remote Island	Goto	25/04/02 - 07/05/02	unknown														
FO-03	N	Fukuokawan	07/05/02 - 17/05/02	70														
FO-05	N	other	10/05/02 - 13/05/02	1														
YG-02	N	other	14/05/02	0.001														
NS-12	remote Island	Goto	17/05/02 - 22/05/02	0.4														
YG-03	N	other	29/05/02 - 05/06/02	unknown														
YG-04	N	other	06/06/02	unknown														
NS-14	remote Island	Goto	10/06/02 - 15/06/02	unknown														
FO-07	N	Fukuokawan	04/07/02 - 11/07/02	70														
SA-06	N	other	05/07/02 - 13/07/02	0.3	Abalone	Turban			died	died		56 kg	130 kg			unknown	unknown	
FO-08	N	Fukuokawan	11/07/02 - 11/08/02	70														
FO-09	N	other	11/07/02 - 02/08/02	1	Abalone				died			unknown				unknown		
SA-07	N	Imariwan	19/07/02 - 22/07/02	5	Amberjacks				died			200 inds				unknown		
NS-17	N	Imariwan	22/07/02	0.015														
NS-20	remote Island	Goto	22/07/02 - 24/07/02	0.0025														
SA-08	N	other	26/07/02 - 28/07/02	0.005														
SA-09	N	Imariwan	26/07/02 - 27/07/02	0.06	Pearl shell				died			5,000 inds.				unknown		
NS-23	N	other	30/07/02 - 31/07/02	0.001														
FO-10	N	Fukuokawan	12/08/02 - 21/08/02	70														
NS-26	remote Island	Goto	24/08/02 - 27/08/02	unknown	Amberjacks	Horse mackerel			died	died		9,280 inds.	620 inds.		29,044	1,240		
NS-27	remote Island	Tsushima	05/09/02 - 13/09/02	0.005														
NS-28	remote Island	Tsushima	06/09/02 - 12/09/02	0.35														
SA-12	N	other	09/09/02 - 14/09/02	0.005														
NS-29	remote Island	Tsushima	10/09/02 - 13/09/02	0.0006														
FO-12	N	Fukuokawan	19/09/02 - 24/09/02	70														
SA-13	N	other	19/09/02 - 03/10/02	6.5														
YG-05	N	other	24/09/02 - 01/10/02	unknown	Yellowtail				died			2,000 inds.				15,000		
NS-23	N	other	09/11/02 - 14/11/02	0.02														
FO-15	N	Fukuokawan	02/11/04	70														
YG-06	N	other	28/11/02	0.001														
NS-36	remote Island	Goto	29/11/02 - 01/12/02	0.07														
SA-17	N	other	09/12/02 - 28/12/02	4.16														
YG-07	N	other	21/12/02	0.005														

Source : Kyushu Fishery Coordination Office, "Situation of Red Tide in the Seas surrounding Kyushu Island, 1999-2003.

Red tide events in Korea (1)

Event No.	Location (name of the sea area)		Duration dd/mm/yy-dd/mm/yy	Continuous days	Causative species				Max. cell density (cells/L)				mitigation activity and effectiveness	Damage	
	Location 1	Location 2												Fisheries resource	Human health
1	Tongyoung buksinman		01-01-99 -		<i>Heterocapsa triquetra</i>					5,200,000					
2	Sachun hangchondong		23-01-99 -		<i>Eutreptiella sp.</i>					3,000,000					
3	Tongyoung hansanman		28-01-99 -		<i>Gymnodinium sp.</i>					780,000					
4	Tongyoung kwangdomyoung		19-04-99 -		<i>Noctiluca sp.</i>					830,000					
5	Masanman sanho		23-04-99 -		<i>Prorocentrum sp.</i>					19,800,000					
6	Geoje ilunmyoung		26-04-99 -		<i>Noctiluca sp.</i>					5,450,000					
7	Masan nanpo		26-04-99 -		<i>Eutreptiella gymnastica</i>					31,000,000					
8	Jinhaeman haengam		28-04-99 -		<i>Eutreptiella gymnastica</i>					16,500,000	1,260				
9	Gunsan naehang		01-05-99 -		<i>Mesodinium rubrum</i>					800,000					
10	Jinhaeman haengam		06-05-99 -		<i>Gymnodinium sanguineum</i>					1,320,000					
11	Ulsan		13-05-99 -		<i>Rhizosolenia sp.</i>					2,000,000	10,000,000				
12	Tongyoung, bubsongman, buksin		15-05-99 -		<i>Heterosigma akashiwo</i>	Proro.minimum	Eutreptiella gymnastica			4,500,000	1,800,000	1,200,000			
13	Masanman		24-05-99 -		<i>Gymnodinium sanguineum</i>					3,960,000					
14	Tongyoung womoonman		24-05-99 -		<i>Mesodinium rubrum</i>					6,500,000					
15	Tongyoung buksinman		28-05-99 -		<i>Leptocylindrus danicus</i>	<i>Gymnodinium sanguineum</i>				7,300,000	1,460,000				
16	kunsan		02-06-99 -		<i>Heterosigma akashiwo</i>					12,000,000					
17	masanman	haengamman	04-06-99 -		<i>Prorocentrum sp.</i>	<i>Heterosigma akashiwo</i>				12,500,000	15,000,000				
18	Namhaegun		08-06-99 -		<i>Ceratium furca</i>					6,000,000					
19	Yeosu gamakmang		09-06-99 -		<i>Heterosigma akashiwo</i>					16,700,000					
20	Pohang youngilman		18-06-99 -		<i>Prorocentrum sp.</i>	<i>Ceratium furca</i>	<i>Heterosigma sp.</i>			650,000	400,000	920,000			
21	Tongyoung kwangdomyoung		18-06-99 -		<i>Prorocentrum sp.</i>					6,600,000					
22	Geoje		19-06-99 -		<i>Prorocentrum sp.</i>	<i>Gymnodinium sp.</i>				330,000	760,000				
23	Tongyoung buksinman		21-06-99 -		<i>Prorocentrum triestinum</i>					5,600,000					
24	Namhae kanginman		21-06-99 -		<i>Prorocentrum sp.</i>					1,350,000					
25	Pusan kadukdo		28-06-99 -		<i>Prorocentrum sp.</i>	<i>Coscinodiscus gigas</i>	<i>Thalassiosira decipiens</i>			15,000,000	1,000,000	100,000			
26	Wando		28-06-99 -		<i>Heterosigma akashiwo</i>					13,000,000					
27	kunsan		29-06-99 -		<i>Heterosigma akashiwo</i>	<i>Heterocapsa triquetra</i>				18,000,000	2,000,000				
28	Pohang youngilman		01-07-99 -		<i>Proro.triquetra</i>	<i>Heterosigma akashiwo</i>	<i>Chaetoceros sp.</i>			600,000	1,500,000	1,600,000	1,500,000		
29	Junnam young		06-07-99 -		<i>Noctiluca scintillans</i>					1,100,000					
30	Tongyoung buksinman		06-07-99 -		<i>Leptocylindrus danicus</i>					2,500,000					
31	Yeosu	Idolsan	06-07-99 -		<i>Skeletonema costatum</i>	<i>Thalassiosira sp.</i>	<i>Ceratium furca</i>	<i>Prorocentrum sp.</i>		2,000,000	5,600,000	250,000	1,700,000		
32	Namhaedo kanginman		07-07-99 -		<i>Chaetoceros sp.</i>					1,600,000					
33	Junnam		08-07-99 -		<i>Chaetoceros sp.</i>	<i>Coscinodiscus gigas</i>	<i>Ceratium furca</i>	<i>Prorocentrum sp.</i>		1,500,000	2,000,000	1,400,000	300,000		
34	Pohang youngilman		21-07-99 -		<i>Prorocentrum triestinum</i>	<i>Prorocentrum micans</i>				6,200,000	1,500,000				
35	Buankun widomyoung		22-07-99 -		<i>Noctiluca scintillans</i>					1,000,000					
36	Jinhaeman	Masan	22-07-99 -		<i>Prorocentrum dentatum</i>	<i>Ceratium furca</i>				4,180,000					
37	Gosung		07-08-99 -		<i>Gymnodinium sp.</i>					15,500,000					
38	Asanman		08-08-99 -		<i>Mesodinium rubrum</i>					4,537,000					
39	Namhaekun	hadongkun	10-08-99 -		<i>Heterosigma sp.</i>	<i>Chaetoceros sp.</i>				2,000,000	3,000,000				
40	Yeosu	Kamakman	10-08-99 -		<i>Chaetoceros sp.</i>	<i>Skeletonema costatum</i>	<i>Nitzschia sp</i>	<i>Rhizosolenia sp.</i>		16,800,000	11,200,000	3,600,000	1,100,000		
41	Ulsan		11-08-99 -		<i>Skeletonema costatum</i>	<i>Chaetoceros sp.</i>				20,000,000	10,000,000				
42	Chunsuman		11-08-99 -		<i>Chaetoceros sp.</i>	<i>Thalassiosira decipiens</i>	<i>Skeletonema costatum</i>	<i>Microcystis viridis</i>		25,000,000	7,500,000	5,000,000	10,000,000		
43	Masan, sanho, dukdong		11-08-99 -		<i>Skeletonema costatum</i>	<i>Rhizosolenia fragilissima</i>				1,760,000	2,840,000				
44	Goheung		11-08-99 - 26/09/99	35	<i>Cochlodinium polykrikoides</i>					260,000					Clay dispersion
45	Yeosu hwajungmyoung		11-08-99 - 22/09/99	34	<i>Cochlodinium polykrikoides</i>					50,000					Clay dispersion
46	Pohang youngilman		13-08-99 -		<i>Prorocentrum sp.</i>	<i>Heterosigma sp.</i>	<i>Skeletonema costatum</i>			2,500,000	1,500,000	2,000,000			
47	Namhaekun sangju		14-08-99 - 17/09/99	33	<i>Cochlodinium polykrikoides</i>					300,000					Clay dispersion
48	Tongyoung		14-08-99 - 01/10/99	45	<i>Cochlodinium polykrikoides</i>					500,000					Clay dispersi finfish died
49	geoje		16-08-99 -		<i>Gymnodinium sanguineum</i>					1,870,000					
50	Ulsan		17-08-99 -		<i>Chaetoceros sp.</i>	<i>Prorocentrum sp.</i>	<i>Thalassiosira sp.</i>			200,000	500,000	200,000			
51	Masan	Jinhaeman	17-08-99 -		<i>Gymnodinium sanguineum</i>	<i>Ceratium sp.</i>				5,280,000	1,760,000				
52	Geoje		18-08-99 -		<i>Gymnodinium sanguineum</i>	<i>Ceratium sp.</i>				1,980,000	3,700,000				
53	Masanman		19-08-99 -		<i>Skeletonema costatum</i>					3,440,000					
54	Jinhaeman		19-08-99 -		<i>Gymnodinium sanguineum</i>					3,080,000					
55	Geoje		21-08-99 -		<i>Gymnodinium mikimotoi</i>					7,500,000					
56	Gosung		21-08-99 - 06/09/99	8	<i>Cochlodinium polykrikoides</i>					2,700,000					Clay dispersion
57	Wando		24-08-99 - 25/09/99	28	<i>Cochlodinium polykrikoides</i>					500,000					Clay dispersion
58	Geoje		25-08-99 - 02/10/99	24	<i>Cochlodinium polykrikoides</i>					1,700,000					Clay dispersion
59	Janghueng		25-08-99 - 20/09/99	22	<i>Cochlodinium polykrikoides</i>					4,000,000					Clay dispersion
60	Pusan		28-08-99 - 03/09/99	5	<i>Cochlodinium polykrikoides</i>					1,100,000					Clay dispersion
61	Ulsan		29-08-99 - 07/09/99	9	<i>Cochlodinium polykrikoides</i>					4,500,000					Clay dispersi finfish died
62	Kyoungju		29-08-99 - 06/09/99	9	<i>Cochlodinium polykrikoides</i>					3,000,000					Clay dispersion
63	Pohang youngilman		31-08-99 - 09/09/99	11	<i>Cochlodinium polykrikoides</i>					3,000,000					Clay dispersion
64	Jinhaeman		31-08-99 - 07/09/99	10	<i>Cochlodinium polykrikoides</i>					7,890,000					Clay dispersion
65	Pohang		02-09-99 - 09/09/99	11	<i>Cochlodinium polykrikoides</i>					3,800,000					
66	Janghueng		06-09-99 -		<i>Gymnodinium sp.</i>					4,300,000					Clay dispersion
67	Namhae		06-09-99 -		<i>Chaetoceros sp.</i>					1,000,000					Clay dispersion
68	Geoje		13-09-99 -		<i>Gymnodinium sanguineum</i>					500,000					
69	Gokun		14-09-99 -		<i>Gymnodinium sp.</i>					2,500,000					
70	Wonmunman		15-09-99 -		<i>Prorocentrum triestinum</i>					3,100,000					

Red tide events in Korea (3)

Event No.	Location (name of the sea area)		Duration dd/mm/yy-dd/mm/yy	Continuous days	Causative species				Max. cell density (cells/L)				mitigation activity and effectiveness	Damage		
	Location 1	Location 2												Fisheries resource	Human health	
141	Kyungbuk		25-08-00		<i>Ceratium furca</i>											
142	Onsanman		28-08-00		<i>Thalassiosira rotula</i>					180,000						
143	Pohang		28-08-00		<i>Chaetoceros sp.</i>	<i>Skeletonema costatum</i>	<i>Ceratium furca</i>			6,800,000	6,000,000	400,000				
144	Ulsan		28-08-00		<i>Prorocystis sp.</i>					40,000						
145	Ulsan		29-08-00		<i>Prorocentrum dentatum</i>	<i>Thalassiosira rotula</i>				84,000	128,000					
146	Yeosu		22-08-00	10/09/00	19	<i>Cochlodinium polykrikoides</i>				910,000						Clay dispersion
147	Tongyoung		24-08-00	11/09/00	19	<i>Cochlodinium polykrikoides</i>				900,000						Clay dispersi finfish died
148	Geoje		02-09-00	2000/9/12	11	<i>Cochlodinium polykrikoides</i>				1,540,000						Clay dispersion
149	Pusan		07-09-00	2000/9/7	1	<i>Cochlodinium polykrikoides</i>				5,000,000						
150	Ulsan		08-09-00			<i>Thalassiosira rotula</i>	<i>Prorocentrum sp.</i>			1,320,000	1,600,000					
151	Ulsan		19-09-00			<i>Ceratium furca</i>				800,000						
152	Geoje		19-09-00			<i>Noctiluca scintillans</i>				450,000						
153	Namhae		20-09-00			<i>Skeletonema costatum</i>				30,000,000						
154	Ulsan		21-09-00			<i>Leptocylindrus danicus</i>	<i>Ceratium furca</i>			16,000,000	53,000					
155	Miinman		22-09-00			<i>Thalassiosira sp.</i>	<i>Mesodinium rubrum</i>	<i>Skeletonema costatum</i>		10,450,000	7,700,000	6,300,000				
156	Ulsan		26-09-00			<i>Thalassiosira decipiens</i>				2,280,000						
157	Ulsan		29-09-00			<i>Prorocentrum sp.</i>				248,000						
158	Geoje		05-10-00			<i>Gymnodinium sanguineum</i>				300,000						Clay dispersion
159	Geoje		08-10-00			<i>Ceratium furca</i>				350,000						
160	Hwado		11-10-00			<i>Gym.sanguineum</i>				850,000						
161	Tongyoung		27-11-00			<i>Proro. Micans</i>				6,200,000						
162	Tongyoung		2001/1/26			<i>Prorocentrum micans</i>				700,000						
163	Pohang		2001/3/19			<i>Cryptomonas acuta</i>				158,400						
164	Pohang		2001/4/7			<i>Eutreptiella gymnastica</i>				1,600,000						
165	Masanman		2001/4/20			<i>Pseudonitschia pungens</i>	<i>Thalassiosira rotula</i>	<i>Eutreptiella gymnastica</i>	<i>Heterosigma akashiwo</i>	1,250	930000	4350	400000			
166	Masanman		2001/5/28			<i>Heterosigma akashiwo</i>				12,350,000						
167	Pusan		2001/5/28			<i>Heterosigma akashiwo</i>				2,800,000						
168	Ulsan		2001/5/28			<i>Heterosigma akashiwo</i>				2,000,000						
169	Pohang		2001/5/29			<i>Heterosigma akashiwo</i>				11,600,000						
170	Onsanman		2001/6/1			<i>Heterosigma akashiwo</i>				7,200,000						
171	Tongyoung		2001/6/1			<i>Prorocentrum micans</i>				1,500,000						
172	Pohang		2001/6/4			<i>Cryptomonas sp.</i>				#####						
173	Ulsan		2001/6/8			<i>Skeletonema costatum</i>	<i>Prorocentrum micans</i>			300,000	1000000					
174	Pohang		2001/6/8			<i>Heterosigma akashiwo</i>				30,000,000						
175	Ulsan		2001/6/8			<i>Prorocentrum triestinum</i>				1,000,000						
176	Pohang		2001/6/13			<i>Eutreptiella gymnastica</i>				40,000,000						
177	Tongyoung		2001/6/21			<i>Heterosigma akashiwo</i>				2,200,000						
178	Tongyoung		2001/6/22			<i>Prorocentrum micans</i>	<i>Prorocentrum triestinum</i>			900,000	1100000					
179	Jindongman		2001/6/22			<i>Heterosigma akashiwo</i>				900,000						
180	Yeosu		2001/6/27			<i>Prorocentrum sp.</i>	<i>Skeletonema costatum</i>			3,600,000	5000000					
181	Masan		2001/7/3			<i>Thalassiosira decipiens</i>	<i>Heterosigma akashiwo</i>			8,900,000	940000					
182	Pusan		2001/7/3			<i>Prorocentrum micans</i>	<i>Skeletonema costatum</i>			1,200,000	9000000					
183	Ulsan		2001/7/3			<i>Heterosigma akashiwo</i>				1,200,000						
184	Jinhaeman		2001/7/3			<i>Thalassiosira decipiens</i>				8,900,000						
185	Boryoung		2001/7/6			<i>Mesodinium rubrum</i>				12,500,000						
186	Suchun		2001/7/9			<i>Mesodinium rubrum</i>	<i>Dictyocha fibula</i>	<i>Cratium sp.</i>	<i>Nitzschia pungens</i>	1,500,000	250000	400000	300000			
187	Masan	Jinhaeman	2001/7/12			<i>Prorocentrum triestinum</i>	<i>Prorocentrum micans</i>			1,240,000	520000					
188	Geoje		2001/7/12			<i>Cratium sp.</i>				1,000,000						
189	Jinhaeman		2001/7/18			<i>Ceratium furca</i>				1,500,000						
190	Kadukdo		2001/7/18			<i>Leptocylindrus danicus</i>	<i>Skeletonema costatum</i>	<i>Nitzschia pungens</i>	<i>Chaetoceros sp.</i>	4,800,000	2000000	2000000	3000000			
191	Kwangyangman		2001/7/18			<i>Heterosigma akashiwo</i>				5,000,000						
192	Dolsando		2001/7/20			<i>Chaetoceros sp.</i>	<i>Skeletonema costatum</i>			2,000,000	6500000					
193	Pusan		2001/7/23			<i>Ceratium furca</i>	<i>Prorocentrum triestinum</i>			350,000	500000					
194	Onsanman		2001/7/24			<i>Leptocylindrus danicus</i>	<i>Chaetoceros sp.</i>	<i>Ceratium furca</i>		5,400,000	3300000	30000				
195	Ulsan		2001/7/24			<i>Prorocentrum triestinum</i>	<i>Leptocylindrus danicus</i>			1,000,000	7800000					
196	Pusan		2001/7/30			<i>Chaetoceros sp.</i>	<i>Prorocentrum sp.</i>			2,000,000	100000					
197	Pusan		2001/7/30			<i>Prorocentrum triestinum</i>				100,000						
198	Pusan		2001/7/30			<i>Skeletonema costatum</i>				1,500,000						
199	Tongyoung		2001/7/30			<i>Prorocentrum triestinum</i>				1,200,000						
200	Ulsan		2001/8/2			<i>Heterosigma akashiwo</i>	<i>Ceratium furca</i>			3,000,000	50000					
201	Onsanman		2001/8/2			<i>Heterosigma akashiwo</i>				8,000,000						
202	Masan		2001/8/4			<i>Prorocentrum micans</i>	<i>Prorocentrum sp.</i>			6,300,000	6500000					
203	Pusan		2001/8/8			<i>Skeletonema costatum</i>	<i>Pseudonitschia pungens</i>			1,400,000	200000					
204	Onsanman		2001/8/10			<i>Prorocentrum minimum</i>				13,000,000						
205	Pohang		2001/8/14			<i>Eutreptiella gymnastica</i>	<i>Heterosigma akashiwo</i>	<i>Prorocentrum triestinum</i>		4,000,000	2000000	3000000				
206	Junnam		2001/8/14	2001/9/9	23	<i>Cochlodinium polykrikoides</i>				600,000						Clay dispersi finfish died
207	Yeosu		2001/8/14	2001/9/8	23	<i>Cochlodinium polykrikoides</i>				9,500,000						Clay dispersi finfish died
208	Namhae		2001/8/15	2001/9/8	18	<i>Cochlodinium polykrikoides</i>				500,000						Clay dispersi finfish died
209	Tongyoung		2001/8/16	2001/9/16	31	<i>Cochlodinium polykrikoides</i>				900,000						Clay dispersi finfish died
210	Uju		2001/8/17			<i>Prorocentrum triestinum</i>				100,000						finfish died

Red tide events in Korea (4)

Event No.	Location (name of the sea area)		Duration		Continuous days	Causative species				Max. cell density (cells/L)				mitigation activity and effectiveness	Damage		
	Location 1	Location 2	dd/mm/yy	-dd/mm/yy											fisheries resource	Human health	
211	Junnam()		2001/8/17	2001/9/11	26	<i>Cochlodinium polykrikoides</i>						410,000			Clay dispersi	finfish died	
212	Pusan		2001/8/18			<i>Prorocentrum triestinum</i>	<i>Skeletonema costatum</i>	<i>Pseudonitschia pungens</i>				100,000	150000	50000			
213	Pusan		2001/8/22			<i>Chaetoceros sp.</i>	<i>Skeletonema costatum</i>	<i>Pseudonitschia pungens</i>				300,000	80000	50000		finfish died	
214	Geoje		2001/8/23	2001/9/12	17	<i>Cochlodinium polykrikoides</i>						680,000			Clay dispersi	finfish died	
215	Pusan		2001/8/24	2001/9/12	20	<i>Cochlodinium polykrikoides</i>						900,000				finfish died	
216	Ulsan		2001/8/25	2001/9/12	19	<i>Cochlodinium polykrikoides</i>						750,000				finfish died	
217	Pohang		2001/8/30	2001/9/12	14	<i>Cochlodinium polykrikoides</i>						1,500,000				finfish died	
218	Donghae		2001/9/5	2001/9/24	17	<i>Cochlodinium polykrikoides</i>						1,700,000				finfish died	
219	Geoje		2001/9/28			<i>Gymnodinium sanguineum</i>						4,780,000					
220	Geoje		#####			<i>Alexandrium sp.</i>						4,780,000					
221	Geoje		#####			<i>Alexandrium sp.</i>						6,200,000					
222	Pohang		2002/2/14			<i>Heterocapsa triquetra</i>						9,000,000					
223	Pohang		2002/2/15			<i>Cryptomonas acuta</i>				3		80,000,000					
224	Masanman	Jinhaeman	2002/5/17			<i>Heterosigma akashiwo</i>			+			8,000,000					
225	Danghangman		2002/5/21			<i>Heterosigma akashiwo</i>				0		5,100,000					
226	Jindongman		2002/5/22			<i>Heterosigma akashiwo</i>						11,200,000					
227	Kadukdo		2002/5/23			<i>Heterosigma akashiwo</i>						2,600,000					
228	Pohang		2002/5/24			<i>Cryptomonas acuta</i>						32,000,000					
229	Pusan		2002/5/24			<i>Heterosigma akashiwo</i>						3,000,000					
230	kunsan		2002/6/4			<i>Leptocylindrus danicus</i>	<i>Chroomonas salina</i>					30,000,000					
231	Masanman		2002/6/15			<i>Thalassiosira decipiens</i>	<i>Eutreptiella gymnastica</i>					2,100,000					
232	Ulsan	Onsanman	2002/6/17			<i>Prorocentrum dentatum</i>	<i>Prorocentrum triestinum</i>					5,000,000					
233	Ulsan		2002/6/20			<i>Prorocentrum triestinum</i>	<i>Prorocentrum dentatum</i>	<i>Heterosigma akashiwo</i>				1,000,000	3000000				
234	Masanman		2002/6/27			<i>Eucamphila zodiacus</i>	<i>Prorocentrum sp.</i>					1,500,000	600000				
235	Masanman		2002/7/10			<i>Leptocylindrus danicus</i>	<i>Rhizosolenia fragilissima</i>					1,500,000	120150000				
236	Wonmunman		2002/7/11			<i>Heterosigma akashiwo</i>						24,000					
237	Kamakman		2002/7/11			<i>Nitzschia pungens</i>	<i>Chaetoceros sp.</i>					510,000	350000				
238	Dolsando		2002/7/11			<i>Skeletonema costatum</i>	<i>Thalassiosira decipiens</i>					1,300,000	850000				
239	Masanman		2002/7/18			<i>Heterosigma akashiwo</i>	<i>Prorocentrum sp.</i>	<i>Thalassiosira decipiens</i>				5,500,000	350000	550000			
240	Masanman	Hangamman	2002/7/24			<i>Prorocentrum sp.</i>						3,200,000					
241	Pusan		2002/7/24			<i>Chaetoceros sp.</i>						20,000,000					
242	Geoje		2002/7/30			<i>Noctiluca scintillans</i>						850,000					
243	Kyoungnam		2002/8/5			<i>Akashiwo sanguinea</i>						2,200,000					
244	Geoje		2002/8/7			<i>Akashiwo sanguinea</i>						9,700,000					
245	Pusan		2002/8/16	2002/9/9	25	<i>Gymnodinium sp.</i>	<i>Cochlodinium polykrikoides</i>					1,100,000					
246	Janheung		2002/8/17			<i>Prorocentrum sp.</i>						280,000					
247	Geoje		2002/8/18	2002/9/11	25	<i>Alexandrium sp.</i>	<i>Cochlodinium polykrikoides</i>					4,200,000			clay dispersi	finfish died	
248	Tongyoung		2002/8/18	2002/9/12	25	<i>Prorocentrum sp.</i>	<i>Cochlodinium polykrikoides</i>					2,300,000			clay dispersi	finfish died	
249	Sachun		2002/8/18	2002/8/24	7	<i>Ceratium furca</i>	<i>Cochlodinium polykrikoides</i>					600,000	300000		clay dispersion		
250	Gosung		2002/8/19	2002/8/31	13	<i>Gymnodinium sanguineum</i>	<i>Cochlodinium polykrikoides</i>					800,000			clay dispersion		
251	Jindongman		2002/8/19			<i>Prorocentrum sp.</i>						2,500,000					
252	Masanman	Hangamman	2002/8/19			<i>Prorocentrum sp.</i>	<i>Thalassiosira decipiens</i>	<i>Skeletonema costatum</i>	<i>Eutreptiella gymnastica</i>			4,500,000	800000	2350000	150000		
253	Namhae		2002/8/19	2002/9/23	35	<i>Skeletonema costatum</i>	<i>Cochlodinium polykrikoides</i>					15,000,000			clay dispersi	finfish died	
254	Pohang		2002/9/20			<i>Chaetoceros sp.</i>	<i>Gymnodinium sanguineum</i>					12,000,000	5000000				
255	Pusan		2002/8/21			<i>Skeletonema costatum</i>						10,000,000					
256	Gosung		2002/8/23			<i>Prorocentrum sp.</i>						1,200,000					
257	Namhae		2002/8/28			<i>Ceratium furca</i>						300,000			clay dispersi	finfish died	
258	Junnam		2002/8/28			<i>Noctiluca scintillans</i>						3,400,000					
259	Jinhae	Hangamman	2002/8/29			<i>Nitzschia pungens</i>	<i>Rhizosolenia fragilissima</i>	<i>Skeletonema costatum</i>				8,700,000	1800000	8700000			
260	Masan	Hangamman	2002/9/5			<i>Prorocentrum sp.</i>						7,500,000					
261	Namhae		2002/9/7			<i>Skeletonema costatum</i>						25,000,000					
262	Namhae		2002/9/9			<i>Skeletonema costatum</i>						15,000,000					
263	Masanman	Jinhaeman	2002/9/10			<i>Skeletonema costatum</i>	<i>Thalassiosira decipiens</i>					3,500,000	6200000				
264	Geoje		2002/10/2			<i>Alexandrium sp.</i>						1,800,000					
265	Pohang		2003/2/7			<i>Cryptomonas acuta</i>						4,000,000					
266	Masanman		2003/4/28			<i>Prorocentrum minimum</i>						32,000,000					
267	Masanman		2003/5/14			<i>Heterosigma akashiwo</i>						32,500,000					
268	Masanman	Hangamman	2003/5/14			<i>Rhizosolenia setigera</i>	<i>Pseudonitschia pungens</i>					4,053,000	2394000				
269	Masanman		2003/5/19			<i>Eucamphila zodiacus</i>						16,650,000					
270	Masanman	Hangamman	2003/5/23			<i>Heterosigma akashiwo</i>						27,800,000					
271	Masanman	Hangamman	2003/6/10			<i>Prorocentrum sp.</i>						3,600,000					
272	Tongyoung		2003/6/12			<i>Akashiwo sanguinea</i>						500,000					
273	Pusan		2003/6/13			<i>Prorocentrum sp.</i>						5,500,000					
274	Kajodo		2003/6/21			<i>Prorocentrum dentatum</i>	<i>Prorocentrum dentatum</i>					2,300,000	2100000				
275	Namhae	Sachun	2003/6/23			<i>Prorocentrum dentatum</i>	<i>Prorocentrum dentatum</i>					4,500,000	35000000				
276	Gosung	Jaranman	2003/6/30			<i>Prorocentrum dentatum</i>	<i>Heterocapsa triquetra</i>	<i>Prorocentrum dentatum</i>				5,100,000	600000	12000000			
277	Kangjinman		2003/7/5			<i>Prorocentrum dentatum</i>	<i>Skeletonema costatum</i>					45,000,000					
278	Tongyoung		2003/7/8			<i>Prorocentrum dentatum</i>						#####					
279	Yeojaman		2003/7/8			<i>Heterosigma akashiwo</i>						20,000,000					
280	Chungnam		2003/7/9			<i>Prorocentrum micans</i>						10,000,000					

Red tide events in Korea (5)

Event No.	Location (name of the sea area)		Duration dd/mm/yy-dd/mm/yy	Continuous days	Causative species					Max. cell density (cells/L)					mitigation activity and effectiveness	Damage		
	Location 1	Location 2														Fisheries resource	Human health	
281	Gosung		2003/7/11		<i>Akashiwo sanguinea</i>					4,500,000								
282	Pohang		2003/7/11		<i>Heterosigma akashiwo</i>					50,000,000								
283	Ulsan		2003/8/5		<i>Chaetoceros sp.</i>					13,000,000								
284	Yeosu		2003/8/13	2003/10/6	47	<i>Cochlodinium polykrikoides</i>					9,500,000				clay dispersi	finfish died		
285	Namhae		2003/8/13	2003/10/7	56	<i>Skeletonema costatum</i>	<i>Cochlodinium polykrikoides</i>				23,000,000	20,000,000			clay dispersi	finfish died		
286	Wando		2003/8/14	2003/10/6	53	<i>Cochlodinium polykrikoides</i>					16,000,000				clay dispersi	abalone died (mass mortality)		
287	Tongyoung		2003/8/14	2003/10/5	52	<i>Cochlodinium polykrikoides</i>					24,000,000				clay dispersi	finfish died		
288	Geoje		2003/8/24	2003/10/6	37	<i>Cochlodinium polykrikoides</i>					7,200,000				clay dispersi	finfish died		
289	Namhae		2003/8/25			<i>Skeletonema costatum</i>					40,000,000							
290	Ulsan		2003/8/27	2003/10/8	42	<i>Cochlodinium polykrikoides</i>					20,000,000							
291	Pohang		2003/8/27	2003/10/8	42	<i>Cochlodinium polykrikoides</i>					16,000,000							
292	Pusan		2003/8/28	2003/10/8	41	<i>Cochlodinium polykrikoides</i>					12,400,000							
293	Ulsan		2003/8/30	2003/10/1	30	<i>Cochlodinium polykrikoides</i>					26,000,000						finfish died	
294	Donghae		2003/9/5	2003/10/5	33	<i>Cochlodinium polykrikoides</i>					23,000,000							
295	Boryoung		2003/9/16			<i>Heterosigma akashiwo</i>					27,000,000							
296	Geoje		2003/9/15			<i>Prorocentrum dentatum</i>					15,600,000							
297	Masanman		2003/9/17			<i>Prorocentrum minimum</i>					30,000,000							
298	Masanman		2003/9/22			<i>Skeletonema costatum</i>					6,150,000							

Red tide events in Russia

Event No.	Location (name of the sea area)	LatitudeN	LongitudeE	Duration dd/mm/yy-dd/mm/yy	Approximate area suffered (km ²)	Type of HAB (Red tide or Toxic)	Causative species	Max. cell density cells/l	Mitigation activity and effectiveness	Damage Fishery resources / Human health
1	Peter the Great Bay	43 11 7	132 16 6	15/06/1992	<1	Red tide	<i>Noctiluca scintillans</i>	450,000	no data	no data
2	Amurskii Bay	43 15 3	131 90 2	25/06/1992	< 1	Red tid	<i>Pseudo-nitzschia pungens/multiseriis</i>	35,000,000	no data	no data
3	Amurskii Bay	43 15 3	131 90 2	15/07/1992	< 1	Red tide	<i>Prorocentrum minimum</i>	8,000,000	no data	no data
4	Peter the Great Bay	43 11 7	132 16 6	05/06/1993	<1	Red tide	<i>Noctiluca scintillans</i>	500,000	no data	no data
5	Amurskii Bay	43 15 3	131 90 2	31/07/1993	< 1	Red tid	<i>Skeletonema costatum</i>	17,400,000	no data	no data
6	Peter the Great Bay	43 11 7	132 16 6	25/05/1994	< 1	Red tide	<i>Noctiluca scintillans</i>	550,000	no data	no data
7	Peter the Great Bay	43 11 7	132 16 6	10/06/1995	< 1	Red tide	<i>Noctiluca scintillans</i>	400,000	no data	no data
8	Amurskii Bay	43 15 3	131 90 2	12/06/1995	< 1	Red tide	<i>Heterosigma akashiwo</i>	5,000,000	no data	no data
9	Amurskii Bay	43 15 3	131 90 2	29/07/1996	< 1	Red tid	<i>Skeletonema costatum</i>	12,700,000	no data	
10	Amurskii Bay	43 15 3	131 90 2	15/07/1997	< 1	Red tide	<i>Skeletonema costatum</i>	3,000,000	no data	no data
11	Amurskii Bay	43 15 3	131 90 2	03/11/1997	< 1	Red tid	<i>Pseudo-nitzschia calliantha/pseudodelicatissima</i>	2,700,000	no data	no data
12	Rynda Bay	43 2 5	131 78 7	11/09/2000	<1	Red tide	<i>Pseudo-nitzschia pungens</i>	1,690,000	no data	no data
13	Rynda Bay	43 2 5	131 78 7	15/08/2000	<1	Red tide	<i>Ditylum brighwellii</i>	1,400,000	no data	no data
14	Golden Horn Bay	43 10 67	131 88 2	12/03/2001	< 1	Red tide	<i>Eutreptia lanowii</i>	15,600,000	no data	no data
15	Golden Horn Bay	43 10 67	131 88 2	10/04/2001	< 1	Red tide	<i>Eutreptiella gymnastica</i>	30,900,000	no data	no data
16	Golden Horn Bay	43 10 67	131 88 2	10/09/2001	< 1	Red tide	<i>Chattonella globosa</i>	6,000,000	no data	no data
17	Rynda Bay	43 2 5	131 78 7	15/05/2002	5	Red tide	<i>Noctiluca scintillans</i>	700,000	no data	no data
18	Amurskii Bay	43 15 3	131 90 2	09/07/2002 - 25/07/2002	< 1	Red tide	<i>Oxyrrhis marina</i>	20,000,000	no data	no data
19	Amurskii Bay	43 15 3	131 90 2	01/08/2002- 06/08/2002	< 1	Red tide	<i>Prorocentrum minimum</i>	11,940,000	no data	no data
20	Amurskii Bay	43 15 3	131 90 2	03/09/2002	< 1	Red tide	<i>Heterosigma akashiwo</i>	7,000,000	no data	no data
21	Vostok Bay	42 88 7	132 72 9	05/05/2003	< 1	Red tide	<i>Noctiluca scintillans</i>	970,000	no data	no data
22	Amurskii Bay	43 15 3	131 90 2	11/05/2003 - 17/06/2003	2	Red tide	<i>Noctiluca scintillans</i>	800,000	no data	no data
23	Amurskii Bay	43 15 3	131 90 2	17/06/2003	< 1	Red tide	<i>Heterosigma akashiwo</i>	25,000,000	no data	no data